

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| (51) International Patent Classification ⁶: C12N 5/06, 5/08 | A1 | (11) International Publication Number: WO 99/57248 (43) International Publication Date: 11 November 1999 (11.11.99) |
| (21) International Application Number: PCT/US98/08716 (22) International Filing Date: 30 April 1998 (30.04.98) (71) Applicant (for all designated States except US): PRESIDENT AND FELLOWS OF HARVARD COLLEGE [US/US]; 17 Quincy Street, Cambridge, MA 02138 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): McMAHON, Andrew, P. [GB/US]; 128 Kendall Road, Lexington, MA 02173 (US). LEE, Scott, K. [US/US]; 54 Chestnut Street, Cambridge, MA 02139 (US). TAKADA, Shinji [-/-]; - (74) Agent: FREEMAN, John, W.; Fish & Richardson P.C., 225 Franklin Street, Boston, MA 02110-2804 (US). | | (81) Designated States: CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> |
| (54) Title: INDUCTION OF NEURONAL REGENERATION (57) Abstract An enriched population of mammalian dorsal neural progenitor cells, e.g., dopaminergic neural precursor cells, are described that are useful to induce neuronal regeneration in mammals suffering from a neurodegenerative disease. | | |

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|------------------------------------------|----|----------------------------------------------|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | | | TR | Turkey |
| BG | Bulgaria | HU | Hungary | ML | Mali | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MN | Mongolia | UA | Ukraine |
| BR | Brazil | IL | Israel | MR | Mauritania | UG | Uganda |
| BY | Belarus | IS | Iceland | MW | Malawi | US | United States of America |
| CA | Canada | IT | Italy | MX | Mexico | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NE | Niger | VN | Viet Nam |
| CG | Congo | KE | Kenya | NL | Netherlands | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NO | Norway | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | NZ | New Zealand | | |
| CM | Cameroon | KR | Republic of Korea | PL | Poland | | |
| CN | China | KZ | Kazakhstan | PT | Portugal | | |
| CU | Cuba | LC | Saint Lucia | RO | Romania | | |
| CZ | Czech Republic | LJ | Liechtenstein | RU | Russian Federation | | |
| DE | Germany | LK | Sri Lanka | SD | Sudan | | |
| DK | Denmark | LR | Liberia | SE | Sweden | | |
| EE | Estonia | | | SG | Singapore | | |

- 1 -

INDUCTION OF NEURONAL REGENERATIONBackground of the Invention

5 The invention relates to neuronal growth and differentiation.

 Wnt polypeptides are secreted cysteine-rich glycosylated polypeptides that play a role in the development of a wide range of organisms. The Wnt family
10 of polypeptides contains at least 16 mammalian members which bind to an extracellular domain of a family of cell surface proteins called Frizzled receptors. Wnt polypeptides may play a role in embryonic induction, generation of cell polarity, and specification of cell
15 fate. Deregulation of Wnt signalling has been linked to tumor development.

Summary of the Invention

 The invention is based on the discovery that Wnt polypeptides regulate neuronal precursor cell fate, i.e.,
20 the type of neuron into which a precursor cell differentiates depends qualitatively on the Wnt signal it receives. For example, Wnt-1 specifies midbrain cell fate. In addition to regulation of cell type, Wnt polypeptides regulate neural precursor state, i.e.,
25 proliferation versus differentiation. A stem cell phenotype is characterized by mitotic activity and a lack of characteristics associated with a mature terminally-differentiated neuron, whereas a differentiated phenotype is characterized by a lack of proliferation and
30 acquisition of properties, e.g., morphology or cell surface proteins, associated with a particular terminally-differentiated neural cell type.

 The invention features an enriched population of mammalian dorsal neural precursor cells that maintain a
35 stem cell phenotype in the presence of a Wnt polypeptide. By an "enriched population" is meant a population of

- 2 -

cells that has been treated with a Wnt polypeptide to selectively expand a desired neural precursor cell type. Thus, an enriched population of neural precursor cells is not naturally-occurring, but contains a higher
5 concentration of neural precursor cells having a particular cell fate compared to the concentration in a naturally-occurring population of stem cells.

The Wnt polypeptide is preferably a Wnt-1 class polypeptide such as Wnt-1, Wnt-2, Wnt-3a, Wnt-7a, and
10 Wnt-7b. A Wnt-1 class polypeptide is a Wnt polypeptide that transforms C57MG cells in culture. Other Wnt polypeptides, e.g., Wnt-5a, that play a role in midbrain development may also be used to culture stem cells.

A drawback of conventional stem cell preparations
15 is that they heterogenous, i.e., a precursor cell with a particular cell fate may constitute only a small fraction of the population. The invention solves this problem by providing a method of selecting for a desired precursor cell type, i.e., by contacting the cell with a Wnt
20 polypeptide. For example, the invention provides a method of treating a heterogeneous population of neural cell precursor cells to enrich for neural precursor cells committed to a desired cell fate by culturing the population with a Wnt polypeptide, e.g., a Wnt-1 class
25 polypeptide. Neural precursor cells selectively proliferate in the presence of the Wnt polypeptide, whereas other precursor cells do not proliferate (or proliferate at a rate lower than that of the dorsal neural precursor cells). Thus, repeated culturing of the
30 population in this manner yields a population of neural precursor cells that is progressively more enriched for dorsal neural precursor cells. The enriched population of dorsal neural precursor cells is at least 60%, preferably at least 75%, more preferably at least 80%,
35 more preferably at least 90%, more preferably at least

- 3 -

95%, more preferably at least 98%, and most preferably 100% dorsal neural precursor cells.

For example, the invention encompasses an enriched population of mammalian dopaminergic neuron precursor
5 cells. Selection of such cells is accomplished by contacting a heterogenous population of precursor cells with a Wnt-1 class polypeptide. The cells may be human or porcine cells and may be derived from fetal tissue. The cells are mitotically-active and maintaining a stem
10 cell phenotype in the presence of a Wnt polypeptide. In the absence of a Wnt polypeptide, the cells cease proliferating and differentiate into dopaminergic neurons. A dopaminergic neuron is one that produces dopamine. Preferably, the Wnt polypeptide is human Wnt-1
15 or a fragment of Wnt-1 that is capable of stimulating proliferation of such cells and arresting differentiation. Since Wnt polypeptides have mitogenic activity for neural precursor cells, a method of stimulating cell proliferation of a dorsal neural
20 precursor cell is carried out by contacting the cell in culture or *in vivo* with a Wnt-1 polypeptide and/or a Wnt-3a polypeptide. To promote proliferation of mammalian dopaminergic neuron precursor cells, the polypeptide preferably has a sequence that is at least 80% identical
25 to amino acid sequence of naturally-occurring human Wnt-1 (SEQ ID NO:1) and has a biological property of naturally-occurring Wnt-1, e.g., the ability to maintain the neural stem cell phenotype of a neural precursor cell in culture.

- 4 -

Table 1: Human Wnt-1 amino acid sequence

1 MGLWALLPGW VSATLLLLALA ALPAALAANS SGRWWGIVNV ASSTNLLTDS
 KSLQLVLEPS
 5 61 LQLLSRKQRR LIRQNPGLIH SVSGGLQSAV RECKWQFRNR RWNCPTAPGP
 HLFQKIVNRG
 121 CRETAFIFAI TSAGVTHSVA RSCSEGSIES CTCDYRRRGP GGPDPWHWGGC
 SDNIDFGRLF
 181 GREFVDSGEK GRDLRFLMNL HNNEAGRTTV FSEMRQECKC HGMSGSCTVR
 TCWMRLPTLR
 10 241 AVGDVLRDRF DGASRVLYGN RGSNRASRAE LLRLEPEDPA HKPPSPHDLV
 YFEKSPNFCT
 301 YSGRLGTAGT AGRACNSSSP ALDGCELLCC GRGHRTRTQR VTERCNCFTF
 WCCHVSCRNC
 361 THTRVLHECL (SEQ ID NO:1)

15 Table 2: Human Wnt-2 amino acid sequence

MNAPLGGIWLWLP LLLTWTLP EVNSSWWYMRATGGSSRV MCDNV
 PGLVSSQRQLCHRHPDVMRAISQGVAEWTAECQHOFQHRWNCNTLDRDHSIFGRVLL
 RSSRESAFVYAISSAGVVFAITRACSQGEVKSCSDPKKMGS AKDSKGI FDWGGCSDN
 IDYGIKFARAFVDAKERK GKDARALMNLHNNRAGRKAVKRFLKQECKCHGVSGSCTLR
 20 TCWLAMADFRKTGDYLWRKYNGAIQVVMNQDGTGFTVANERFKKPTKNDLVYFENSPD
 YCIRDREAGSLGTAGRV CNLTSRGMDSCVMCCGRGYDTSHTVTRMTKCGCKPHWCCA
 RCQDCLEALDVHTCKAPKNADWTTAT (SEQ ID NO:2)

Where a particular polypeptide or nucleic acid
 molecule is said to have a specific percent identity to a
 25 reference polypeptide or nucleic acid molecule of a
 defined length, the percent identity is relative to the
 reference polypeptide or nucleic acid molecule. Thus, a
 peptide that is 50% identical to a reference polypeptide
 that is 100 amino acids long can be a 50 amino acid
 30 polypeptide that is completely identical to a 50 amino
 acid long portion of the reference polypeptide. It might
 also be a 100 amino acid long polypeptide which is 50%
 identical to the reference polypeptide over its entire
 length. In the case of polypeptide sequences which are
 35 less than 100% identical to a reference sequence, the
 non-identical positions are preferably, but not
 necessarily, conservative substitutions for the reference
 sequence. Conservative substitutions typically include
 substitutions within the following groups: glycine and
 40 alanine; valine, isoleucine, and leucine; aspartic acid
 and glutamic acid; asparagine and glutamine; serine and
 threonine; lysine and arginine; and phenylalanine and
 tyrosine.

- 5 -

Sequence identity can be measured using sequence analysis software (for example, the Sequence Analysis Software Package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison, WI 53705), with the default parameters as specified therein.

An enriched population of mammalian dorsal hindbrain precursor cells is also within the invention. Such cells are selected by contacting a heterogenous population of cells with a mixture of a Wnt-1 polypeptide and a Wnt-3a polypeptide. An enriched population of mitotically-active mammalian hippocampal neuron precursor cells are selected by culturing the cells in the presence of a Wnt-1 class polypeptide such as Wnt-3a; the cells maintain a stem cell phenotype in culture in the presence of a Wnt-3a polypeptide. Such precursor cells cease proliferating and differentiate into hippocampal neurons in the absence of the Wnt-3a polypeptide. Preferably, the Wnt-3a polypeptide has a sequence that is at least 80% identical to SEQ ID NO:2 and has a biological property of naturally-occurring Wnt-3a, e.g., the ability to maintain a neural stem cell phenotype in culture.

Table 3: Murine Wnt-3a amino acid sequence

MAPLGYYLLVLCSLKQALGSYPIWWSLAVGPQYSSLSTQPILCAS
 25 IPGLVPKQLRFCRNYVEIMPSVAEGVKAGIQEQHQFRGRRWNCTTVSNSLAIFGPVL
 DKATRESAFVHAIASAGVAFVTRSCAEGSAAICGSSRLQGSPGEGWKWGGCSEDIE
 FGGMVSREFADARENRPDARSAMNRHNNEAGRQAIASHMHLKCKCHGLSGSCEVKTWCW
 WSQPDFRTIGDFLKDKYDSASEMVVEKHRESRGWVETLRPRYTYFKVPTERDLVYYEA
 30 SPNFCEPNPETGSFGTRDTCNVSSHGIDGCDLLCCGRGHNARTERRREKCHCVFHWK
 CYVSCQECTRVYDVHTCK (SEQ ID NO:3)

Table 10: Human Wnt-3a amino acid sequence

CKCHGLSGSC EVKTCWWSQP DFRAIGDFLK DKYDSASEMV VEKHRESRGW
 VETLRPRYTY FKVPTERDLV YYEASPNFCE PNPETGSFGT RDTCNVSSH
 35 GIDGCDLLCC GRGHNARAER RREKRCVVFH WCC (SEQ ID NO:10)

Table 4: Human Wnt-7a amino acid sequence

1 MNRKALRCLG HLFLSLGMVC LRIGGFSSV ALGATIICNK IPGLAPRQRA ICQSRPDAII
 61 VIGEGSQMGL DECQFQFRNG RWNCALGER TVFGKELKVG SRDGAFTYAI IAAGVAHAIT
 121 AACTHGNLSD CGCDKEKQOQ YHRDEGWKWG GCSADIRYGI GFAKVFDAR EIKONARTLM
 181 NLHNNNEAGRK ILEENMKLEC KCHGVSGSCT TKTCWTTLPQ FRELGYVLKD KYNEAVHVEP
 40 241 VRASRNKRPT FLKIKKPLSY RKPMDTDLVY IEKSPNYCEE DPVTGSGVGTQ GRACNKTAPQ

- 6 -

301 ASGCDLMCCG RGYNTHQYAR VWQCNCXFWH CCYVKCNTCS ERTEMYTCK

Table 5: Human Wnt-7b partial amino acid sequence

1 GVSGSCTTKT CWTTLPKFRE VGHLLKEKYN AAVQVEVVRA SRLRQPTFLR IKOLRSYQKP
 61 METDLVYIEK SPNYCEEDAA TGSVGTQGR I CNRTSPGADG CDTMCCGRGY NTHQYTKVWQ
 5 121 CNCK (SEQ ID NO:5)

Table 6: Human Wnt-5a amino acid sequence

1 MAGSAMSSKF FLVALAIFFS FAQVVIEANS WWSLGMNNPV QMSEVYIIGA QPLCSQLAGL
 61 SQGQKKLCHL YQDHMQYIGE GAKTGIKECQ YQFRHRRWNC STVDNTSVFG RVMQIGSRET
 121 AFTYAVSAAG VVNAMSRACR EGELSTCGCS RAARPKDLPR DWLWGGCGDN IDYGYRFAKE
 10 181 FVDARERERI HAKGSYESAR ILMNLHNEA GRRTVYNLAD VACKCHGVSG SCSLKTCWLQ
 241 LADFRKVGDA LKEYDSAAA MRLNSRGKLV QVNSRFNSPT TQDLVYIDPS PDYCVRNEST
 301 GSLGTQGRLC NKTSEGMDGC ELMCCGRGYD QFKTVQTERC HCKFHWCCYV KCKKCTEIVD
 361 QFVCK (SEQ ID NO:6)

Other patterning signals, e.g., Bmp polypeptides
 15 or Hedgehog polypeptides, are also used to induce
 differentiation of an enriched population of neural
 precursor cells into a differentiated neural cell type.

An population of neural precursor cells that is
 enriched for a particular type of precursor cell is
 20 useful clinically, e.g., to repopulate a depleted
 population of a particular type of neuron. Depletion of
 subpopulations of neurons may be the result of the
 progression of a neurodegenerative disease such as
 Parkinson's Disease, Amyotrophic Lateral Sclerosis,
 25 Diffuse Lewy Body Disease, Cortical-basal Ganglionic
 Degeneration, Hallervorden-Spatz Disease, or Myoclonic
 Epilepsy. A method of inducing neuronal regeneration in
 an adult mammal suffering from a neurodegenerative
 disorder is carried out by transplanting into the
 30 affected mammal an enriched population of dorsal neural
 precursor cells such as that cultured in the presence of
 one or more Wnt polypeptides. To promote proliferation
 of the transplanted stem cells *in vivo*, the method may
 also include a step of administering to the mammal a Wnt
 35 polypeptide or Wnt agonist systemically or locally at the
 site of transplantation. For example, a patient
 suffering from Parkinson's disease is treated by
 transplanting into the brain of the patient an enriched

- 7 -

population of dopaminergic neuron precursor cells. A Wnt-1 polypeptide may be administered concurrently or subsequent to transplantation.

The invention also includes a transgenic non-human mammal, e.g., a rodent such as a mouse, the germ cells and somatic cells of which contain a null mutation, e.g., a deletion, in DNA encoding a Wnt polypeptide. These animals can serve as useful models of neural development. By "null mutation" is meant an alteration in the nucleotide sequence that renders the gene incapable of expressing a functional protein product. The mutation could be in a Wnt gene regulatory region or in the coding sequence. It can, e.g., introduce a stop codon that results in production of a truncated, inactive gene product or it can be a deletion of all or a substantial portion of the coding sequence.

Other features and advantages of the invention will be apparent from the following description of the preferred embodiments thereof, and from the claims.

Detailed Description

The invention provides methods of selecting for neural precursor cells that will differentiate into a particular type of neuron upon exposure to a differentiation-inducing condition or composition and methods for growing such precursor cells in culture. The methods permit expansion of precursor cells of a desired cell fate to achieve large number of cells suitable for clinical transplantation.

Neural Stem Cells

Primary neural progenitor cells are obtained from a mammalian source, e.g., fetal CNS precursor tissue such as developing neural crest tissue, using known methods. Such primary cells are cultured in the presence of a Wnt polypeptide such as Wnt-1 class polypeptide (purified from a natural source or produced recombinantly) in

- 8 -

conventional tissue culture medium such as Dulbecco's Modified Eagles Medium (DMEM) containing fetal calf serum or in serum-free tissue culture medium.

Wnt polypeptides regulate neuronal precursor cell fate as well as neural precursor state. Wnt polypeptides that belong to the Wnt-1 class of Wnt polypeptides are preferably used to culture neural precursor cells for the purpose of maintaining a stem cell phenotype and promoting proliferation. A Wnt-1 class polypeptide is a Wnt polypeptide and that transforms C57MG cells in culture. To determine whether a Wnt polypeptide is a Wnt-1 class polypeptide, C57MG cells (an epithelial cell line derived from normal mouse mammary tissue) are cultured in the presence and absence of the Wnt polypeptide using known methods, e.g., that described by Wong et al., 1994, Mol. Cell. Biol. 14:6278-6286, and their morphology assessed for a transformed phenotype. Normal C57MG cells grow in a monolayer with a regular, cuboidal appearance at confluence, whereas culturing C57MG cells in the presence of a Wnt-1 class polypeptide causes the cells to become transformed, i.e., refractile and elongated, growing over other cells in a disorganized manner. Wnt polypeptides of the Wnt-1 class cause C57MG cells to assume a transformed phenotype. Human Wnt polypeptides which belong to the Wnt-1 class include Wnt-1 (GENBANK Accession #139743, Wnt-2 (GENBANK Accession #139750), Wnt-3a, Wnt-7a (GENBANK Accession #2501663), and Wnt-7b (GENBANK Accession #546573). A Wnt polypeptide, e.g., human Wnt-5a (GENBANK Accession #731157), that is not a member of the Wnt-1 class may also be used (with or without a Wnt-1 class polypeptide) to culture neural precursor cells.

The cells are cultured in the presence or absence of feeder cells. Feeder cells may be engineered to produce a recombinant Wnt-1 class polypeptide such as

- 9 -

Wnt-1 and/or Wnt-3a, e.g., by introducing DNA encoding a Wnt polypeptide, e.g., DNA encoding Wnt-1, Wnt-2, Wnt-3a, Wnt-7a or Wnt-7b, into the cell and culturing the cell under conditions that permit expression of the

- 5 recombinant polypeptide and secretion of the polypeptide into the extracellular environment. For example, feeder cells can be transfected with an expression vector containing DNA having the sequence of naturally-occurring Wnt-1, Wnt-2, or Wnt-3a.

10 Table 7: Human Wnt-1 Nucleotide Sequence

```

      1 atgtatgtat gtatgtatgt atgtatgtat acgtgcgtgc acctgtgtgt
gcttgggtgtc
      61 agtggggctc agacatcacc tgattccctg gaactggagt tacagggtggc
tataagccac
    15      121 cacttgggtg ctgagaacag agtccgggcc tctggcagag cagtcagtgc
ttttagccac
      181 tgagccactc tcatcccccc aattatgttc atcttgagtt gggcaggtac
gggtggcgga
      241 taggcctgta atcccagcag tcaactggacc atcatggggt ctacatatta
    20 aacctttatg
      301 ttaggtaggg tcacacagca agatccggtc acaaaaccag caacaacaaa
aaccaaaagg
      361 agccagcttc ttcccacaag cattctttcc ctcagggtctt cagctccatc
tgacagctac
    25      421 tcggctgggt gtccctatct ttctgagcct agttgccaga gaaacaagcc
cggttcatct
      481 tcattgactag cacatctaata gataagcaca ggttgactca aggtgccata
gagtgacact
      541 aggtaccagc agcgacagaa tgacacctat gagtgcacgt cgtaatacac
    30 aaacacacac
      601 acacacacac acacacacac acacacacac tcatgcaccc acctgcaaac
acaattgcag
      661 ccttctggac gtctcctgtc acagcccac ctccttctctg atacactgcg
ttaagtgggtg
    35      721 actgtaacaa aatgacttca tgctctccct gtcttgagcc aaattacaca
attatttga
      781 aagggctcaa aatgtttctt gttagaagtt tctggataca ccaatacaca
ggagcgtgca
      841 ccctcagaac acatgtacac tttgacttaa tctcacgggt gacacaccga
    40 cgcttacact
      901 cccctagcc cacagaggca aactgctggg cgcttctgag tttctcactg
ccaccagctc
      961 ggtttgtcga gcctaccccc gcaccccgcg cccgggaatc cctgaccaca
gctccaccca
    45      1021 tgctctgtct ccttcttttc cttctctgtc cagccgtcgg ggttcctggg
tgaggaagtg
      1081 tctccacgga gtcgctggct agaaccacaa ctttcatcct gccattcaga
ataggggaaga
      1141 gaagagacca cagcgtaggg gggacagagg agacggactt cgagaggaca
    50 gccccaccgg
      1201 cgcgtgtggg ggaggcaatc caggctgcaa acaggttgct cccagcgcgt
tgtccccgcg
      1261 cccctggcg gatgctgggt cccgacgggc tccggacgcg cagaagagtg
aggccggcgc
    55      1321 gcgtgggagg ccatcccaag gggaggggtc ggcggccagt gcagacctgg
aggcggggcc

```

- 10 -

1381 accaggcagg gggcgggggt gagccccgac ggtagcctg tcagctcttt
 gctcagaccg
 1441 gcaagagcca cagcttcgct cgccactcat tgtctgtggc cctgaccagt
 5 gcgccttggg
 1501 gcttttagtg ccgcccgggc ccggaggggc agcctcttct cactgcagtc
 agcgccgcaa
 1561 ctataagagg cctataagag gcggtgcctc ccgcagtggc tgcttcagcc
 cagcagccag
 1621 gacagcgaac catgctgcct gcggcccgc tccagactta ttagagccag
 10 cctgggaact
 1681 cgcactcactg ccctcaccgc tgtgtccagt cccaccgtcg cggacagcaa
 ccacagtcgt
 1741 cagaaccgca gcacagaacc agcaaggcca ggcaggccat ggggctctgg
 gcgctgctgc
 1801 ccagctgggt ttctactacg ttgctactgg cactgaccgc tctgcccgca
 15 gccctggctg
 1861 ccaacagtag tggccgatgg tggaagtga gctagtacgg ggtccgccac
 ttgtcctggg
 1921 gcaaagagcc aggcacgggc cttaccacgc tcccacgctg tggggatcac
 20 caacctacag
 1981 acccccctcg tgcattgtga cttcacatcc aggggtgctca cacctagaac
 tagctctgct
 2041 gaagtggggc acatcattgg catgcagaag cccagatata ccaggctcag
 agaccattcc
 2101 catttaatac gaccccgttt ctgctgagca acagggtcca acctcgctgt
 25 ggtgggtgct
 2161 cagggtgtccc ttaggtcttg aaccaaaaaa aaaaaaaaaa aaaaaaaaaa
 accagatatt
 2221 agctttgagg tgagggagtg gaattcctaa gtttttcaag gtgggcaagg
 30 ctgcagggtg
 2281 ggttttctct cgggggctga cttgaagaaa ggaagagcta aggtagccat
 gccttttctg
 2341 tccactcact agactctgga gctcaggggc aggcaggat aggggtgtac
 agcctgtatg
 2401 gttaggatgc aggtcccctc ccctggactg aacccttatg catcccgcc
 35 ggggcatcgt
 2461 gaacatagcc tcctccacga acctgttgac ggattccaag agtctgcagc
 tgggtgctga
 2521 gccagctctg cagctgctga gccgcaagca gcggcgactg atccgacaga
 40 acccggggat
 2581 cctgcacagc gtgagtggag ggctccagag cgctgtgcga gagtgcaaat
 ggcaattccg
 2641 aaaccgccgc tggaaactgcc ccactgctcc ggggccccac ctcttcggca
 agatcgtcaa
 2701 ccgagggtggg tgcccaggaa agcgacgctt ccgggattaa gggaaaagca
 45 gggcatctc
 2761 cagggcatag gcgggcgaag gcagggaaga catcccaggg ttatatgtga
 tcaaactgag
 2821 aatcgctggt tgccggcagt taccgtagggt cagcaccaga ttctttctag
 50 ccttgcggtg
 2881 tgagcatgat ctttaacggt gctggccact ggcccacaga aagggaaatc
 cggatcgtgg
 2941 gcgctgggag acagctgttt ttccctagcc ttccctcaaag gtacctggga
 agctgatctc
 3001 tgagggttag ctagggtgtg gcttcgcacc cagcaaagtt tgcactgcca
 55 atactagtag
 3061 cgatcttggc tatgcagatt tgttctactt gggaaatctc ccttggagct
 gctctgctag
 3121 ggctctggag tctcagtaaa gcttagagag gagggcatte catgcttcgc
 60 acacatgact
 3181 ccaaggatgt tggactgtag ggtaccaagt cttccaaaca ggggtgctgag
 ttggccccac
 3241 gcctttctct aactgatgcg gggtegettc acccacaggc tgccgagaaa
 cagcgttcat
 3301 ctctcgcaatc acctccgccg gggteacaca ttccgtggcg cgctcctgct
 65 ccgaaggctc

- 11 -

3361 catcgagtc tgcacctgcg actaccggcg gcgcggccct gggggcccg
 actggcactg
 3421 ggggggctgc agtgacaaca tcgatttttg tcgcctcttt ggccgagagt
 tcgtggactc
 5 3481 cggggagaag gggcgggacc tacgcttcct catgaacctt cacaacaacg
 aggcaggcg
 3541 aacggtacgt cgggtgtgtcc ggaaccaatg gcaggggaga tgtaagacag
 gtgcacgggg
 3601 acagaggcac agggaggggc tccccgagag agtgggactc taggagggaa
 10 gacagagaag
 3661 aggtgggtggg tgagggcaaa gaggttcctg agctgatgac agaacagaag
 agattagcag
 3721 gctatcaaca cgtgggatgt attgagatgg ctccatggca cacttttgaa
 agataaaaagt
 15 3781 gacttgctgg cgtggagcag agtctggccg aatgtcccta tctcagcggg
 ccattttgca
 3841 ctctctctct cccgagctta gtcacacctg gaccttggtc gaagtttcca
 cagcatcgac
 3901 gtgacccggg tgggggtggg gtggggaagt atgggtggtg gttcgtggga
 20 tgttggtctt
 3961 gaccttttct tccctctctc cctcgtcccc tctccccca gaccgtgttc
 tctgagatgc
 4021 gccaaagagt caaatgccac gggatgtccg gctcctgcac ggtgcgcacg
 tgttggtatgc
 25 4081 ggctgcccac gctgcgcgct gtgggcgacg tgctgcgcga ccgcttcgac
 ggcgccctcc
 4141 gcgtccttta cggcaaccga ggcagcaacc gcgcctcgcg ggcggagctg
 ctgcgcctgg
 4201 agcccgaaga ccccgcgcac aagcctccct cccctcacga cctcgtctac
 30 ttcgagaaat
 4261 cgcccaactt ctgcacgtac agtggccgccc tgggcacagc tggcacagct
 ggacgagctt
 4321 gcaacagctc gtctcccgcg ctggacggct gtgagctgct gtgctgtggc
 cgaggccacc
 35 4381 gcacgcgcac gcagcgcgct acggagcgtc gcaactgcac cttccactgg
 tgctgccacg
 4441 tcagctgccg caactgcacg cacacgcgcg ttctgcacga gtgtctatga
 ggtgccgcgc
 4501 ctccgggaac gggaacgctc tcttcagtt ctacagacaca ctgcgtggtc
 40 ctgatgtttg
 4561 cccaccctac cgcgtccagc cacagtccca gggttcatag cgatccatct
 ctcccacctc
 4621 ctacctgggg actcctgaaa ccacttgctc gagtcggctc gaaccctttt
 gccatcctga
 45 4681 gggccctgac ccagcctacc tccctccctc tttgagggag actccttttg
 cactgcccc
 4741 caatttggtc agaggggtgag agaaagattc ttcttctggg gtgggggtgg
 ggaggtcaac
 4801 tcttgaaggt gttgcgggtc ctgatgtatt ttgcgctgtg acctcttttg
 50 gtattatcac
 4861 ctttctctgt ctctcgggtc cctataggtc ccttgagttc tctaaccagc
 acctctgggc
 4921 ttcaaggcct tccccctccc acctgtagct gaagagtttc cgagttgaaa
 gggcacggaa
 55 4981 agctaagtgg gaaaggaggt tgctggacct agcagcaaaa ccctacattc
 tccttgctc
 5041 tgcctcgag ccattgaaca gctgtgaacc atgcctccct cagcctcctc
 ccacccttc
 5101 ctgtcctgcc tctcatcac tgtgtaaata atttgcaccg aaatgtggcc
 60 gcagagccac
 5161 gcgttcgggt atgtaaataa aactatztat tgtgctgggt tccagcctgg
 gttgcagaga
 5221 ccaccctcac cccacctcac tgctcctctg ttctgctcgc cagtcctttt
 gttatccgac
 65 5281 cttttttctc ttttaccag cttctcatag gcgccttgc ccaccggatc
 agtatttct

- 12 -

5341 tccactgtag ctattagtgg ctctcgccc ccaccaatgt agtatcttcc
 tctgaggaat
 5401 aaaatatcta tttttatcaa cgactctggt ccttgaatcc agaacacagc
 atggcttcca
 5 5461 acgtcctctt cccttccaat ggacttgctt ctcttctcat agccaaacaa
 aagagataga
 5521 gttgttgaag atctcttttc cagggcctga gcaaggaccc tgagatcctg
 acccttggat
 5581 gaccctaaat gagaccaact agggatc (SEQ ID NO:7)

10 Table 8: Human Wnt-2 Nucleotide Sequence

1 agcagagcgg acgggagcgc gggaggcgcg cagagctttc gggctgcagg cgctcgctgc
 61 cgctggggaa ttgggctgtg ggcgaggcgc tccgggctgg cctttatcgc tcgctggggc
 121 catcgtttga aactttatca gcgagtcgcc actcgtcgca ggaccgagcg gggggcgggg
 181 gcgcggcgag gcggcgggcg tgacgaggcg ctcccgagc tgagcgcttc tgctctgggc
 15 241 acgcatggcg ccgcacacg gagtctgacc tgatgcagac gcaagggggt taatatgaac
 301 gccctctcgc gtggaatctg gctctggctc cctctgctct tgacctggct caccgccgag
 361 gtcaactctt catggtggta catgagagct acaggtggct cctccagggt gatgtgcat
 421 aatgtgccag gcctggtag gagccagcgg cagctgtgtc accgacatcc agatgtgatg
 481 cgtgccatta gccagggcgt ggccgagtg acagcagaat gccagcacca gttccgccag
 20 541 caccgctgga attgcaacac cctggacagg gatcacagcc tttttggcag ggtcctactc
 601 cgaagtagtc ggggaatctgc ctttgtttat gccatctcct cagctggagt tgtatttggc
 661 atcaccaggg cctgtagcca aggagaagta aaatcctgtt cctgtgatcc aaagaagatg
 721 ggaagcgcca aggacagcaa aggcattttt gattgggggt gctgcagtga taacattgac
 781 tatgggatca aatttgcccg cgcattttgt gatgcaagg aaaggaaagg aaaggatgcc
 25 841 agagccctga tgaatcttca caacaacaga gctggcagga aggtgtaaa gcggttcttg
 901 aaacaagagt gcaagtgcc cggggtgagc ggctcatgta ctctcaggac atgctggctg
 961 gccatggccg acttcaggaa aacgggagat tatctctgga ggaagtacaa tggggccatc
 1021 caggtggtca tgaaccagga tggcacaggt ttactgtgg ctaacgagag gtttaagaag
 1081 ccaacgaaaa atgacctcgt gtattttgag aattctccag actactgtat cagggaccga
 30 1141 gaggcaggct ccctgggtac agcaggcgt gtgtgcaacc tgacttccc gggcatggac
 1201 agctgtgaag tcatgtgtc tgaggagcgc tacgacacct cccatgtcac ccgatgacc
 1261 aagtgtgggt gtaagtcca ctggtgtgc gccgtgcgct gtcaggactg cctggaagct
 1321 ctggatgtgc acacatgcaa ggcctccaa aacgctgact ggacaaccgc tacatgacct
 1381 cagcaggcgt caccatccac ctccctctc acaaggactc cattggatct gcaagaacac
 35 1441 tggacctttg gggtctttct ggggggatat ttcttaaggc atgtggcctt tatctcaacg
 1501 gaagccccct ctctctccct gggggcccca ggaagggggg ccacacgctg cacctaaagc
 1561 ctacctatt ctatccatct cctggtgttc tgcagtcac tccctcctg gcgagttctc
 1621 tttggaaata gcatgacagg ctgttcagcg gggagggtgg tggggccaga ccactgtctc
 1681 caccacctt gacgtttctt ctttctagag cagttggcca agcagaaaaa aaagtgtctc
 40 1741 aaaggagctt tctcaatgtc ttcccaaaa tgggtccaat taagaaattc catacttctc
 1801 tcagatggaa cagtaaagaa agcagaatca actgcccctg acttaacttt aacttttgaa
 1861 aagaccaaga cttttgtctg tacaagtgg tttacagcta ccacccttag ggttaattgt
 1921 aattacctgg agaagaatgg ctttcaatac ccttttaagt ttaaaatgtg tatttttcaa
 1981 ggcatttatt gccatattaa aatctgatgt aacaagggtg ggacgtgtgt cctttggtac
 45 2041 tatggtgtgt tgtatctttg taagagcaaa agcctcagaa agggattgct ttgcattact
 2101 gtcccttga tataaaaaat ctttaggaa tgagagttcc ttctcactta gaatctgaag
 2161 ggaattaaaa agaagatgaa tggctcggca atattctgta actattgggt gaatatgggt
 2221 gaaaataatt tagtggatgg aatatcagaa gtatatctgt acagatcaag aaaaaaggga
 2281 agaataaaat tcctatatca t (SEQ ID NO:8)

50 Table 9: Murine Wnt-3A Nucleotide Sequence

1 gaattcatgt cttacgggtca aggcagaggg ccagcgcga ctgcagccgc
 gccacctccc
 61 agggccgggc cagcccaggc gtccgcgctc tcgggggtgga ctccccccgc
 55 tgcgcgctca
 121 agccggcgat ggctcctctc ggatacctct tagtgctctg cagectgaag
 caggctctgg
 181 gcagctaccc gatctgggtg tcttggctg tgggacccca gtactcctc
 ctgagcactc
 60 241 agccattct ctgtgccagc atcccaggcc tggtagcgaa gcagctgcgc
 ttctgcagga

- 13 -

301 actacgtgga gatcatgccc agcgtggctg aggggtgtcaa agcgggcatc
 caggagtggc
 361 agcaccagtt ccgaggccgg cgttggaact gcaccaccgt cagcaacagc
 ctggccatct
 5 421 ttggccctgt tctggacaaa gccacccggg agtcagcctt tgtccatgcc
 atcgccctccg
 481 ctggagtagc ttctgcagtg acacgctcct gtgcagaggg atcagctgct
 atctgtgggt
 541 gcagcagccg cctccagggc tccccaggcg agggctggaa gtggggcggc
 10 tgtagtgagg
 601 acattgaatt tggaggaatg gtctctcggg agtttgccga tgccagggag
 aaccggccgg
 661 atgcccgtc tgccatgaac cgtcacaaca atgaggctgg gcgccaggcc
 atcgccagtc
 15 721 acatgcacct caagtgc aaa tgccacgggc tatctggcag ctgtgaagt
 aagacctgct
 781 ggtggtcgca gccggacttc cgcaccatcg gggatttct caaggacaag
 tatgacagt
 841 cctcggagat ggtggttagag aaacaccgag agtctcgtgg ctgggtggag
 20 accctgaggc
 901 cacgttacac gtacttcaag gtgccgacag aacgcgacct ggtctactac
 gaggccctcac
 961 ccaacttctg cgaacctaac cccgaaaccg gtccttcgg gacgcgtgac
 cgcacctgca
 1021 atgtgagctc gcatggcata gatgggtgcg acctgttgtg ctgcgggcgc
 25 gggcataacg
 1081 cgcgcactga gcgacggagg gagaaatgcc actgtgtttt ccattggtgc
 tgctacgtca
 1141 gctgccagga gtgcacacgt gtctatgacg tgcacacctg caagtaggag
 30 agctcctaac
 1201 acgggagcag ggttcattcc gaggggcaag gttcctacct gggggcggg
 ttctacttg
 1261 gaggggtctc ttacttgggg actcggttct tacttgaggg cggagatcct
 acctgtgagg
 1321 gtctcatacc taaggacctg gtttctgcct tcagcctggg ctctatttg
 35 ggatctgggt
 1381 tccttttttag gggagaagct cctgtctggg atacgggttt ctgcccagg
 gtggggctcc
 1441 acttggggat ggaattccaa tttgggccgg aagtcctacc tcaatggctt
 40 ggactcctct
 1501 cttgaccga cagggtcaa atggagacag gtaagctact ccctcaacta
 ggtgggggtc
 1561 gtgcggatgg gtgggagggg agagattagg gtccctcctc ccagaggcac
 tgctctatct
 1621 agatacatga gaggggtgctt cagggtgggc cctatttggg cttgaggatc
 45 ccgtgggggc
 1681 ggggcttcac cccgactggg tggaaactttt ggagaccccc ttccactggg
 gcaaggcttc
 1741 actgaagact catgggatgg agctccacgg aaggaggagt tctgagcga
 50 gcctgggctc
 1801 tgagcaggcc atccagctcc catctggccc cttccagtc ctggtgtaag
 gttcaacctg
 1861 caagcctcat ctgcgcagag caggatctcc tggcagaatg aggcattggag
 aagaactcag
 1921 gggatgatac aagacctaac aaaccccggt cctgggtacc tcttttaaag
 55 ctctgcaccc
 1981 cttcttcaag ggctttccta gtctccttgg cagagctttc ctgaggaaga
 tttgcagtcc
 2041 cccagagtcc aagtgaacac ccatagaaca gaacagactc tatcctgagt
 60 agagaggggt
 2101 ctctaggaat ctctatgggg actgctagga aggatcctgg gcatgacagc
 ctctgtatgat
 2161 agcctgcac cgtctgaca cttaatactc agatctccc ggaacccag
 ctcatccgt
 2221 ccgtgatgtc catgccccaa atgcctcaga gatgttgctt cactttgagt
 65 tgtatgaact

- 14 -

```

      2281 tcggagacat ggggacacag tcaagccgca gagccagggt tgtttcagga
cccatctgat
      2341 tccccagagc ctgctgttga ggcaatgggc accagatccg ttggccacca
ccctgtcccg
5      2401 agcttctcta gtgtctgtct ggcttgaag tgagggtgcta catacagccc
atctgccaca
      2461 agagcttctt gatttggtacc actgtgaacc gtccctcccc ctccagacag
gggaggggat
10     2521 gtggccatac aggagtgtgc ccggagagcg cggaaagagg aagagagggt
gcacacgcgt
      2581 ggtgactgac tgtcttctgc ctggaacttt gcgttcgcgc ttgtaacttt
attttcaatg
      2641 ctgctatata caccacaccac tggatttaga caaaagtgat tttctttttt
tttttttctt
15     2701 ttctttctat gaaagaaatt attttagttt atagtatgtt tgtttcaaat
aatggggaaa
      2761 gtaaaaagag agaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaa
      (SEQ ID NO:9)

```

Table 11: Human Wnt-3a nucleotide sequence

```

20     tgtaagtgcc acgggctgtc gggcagctgc gaggtgaaga catgctgggtg
      gtcgcaaccc gacttccgcg ccatcgggtga cttcctcaag gacaagtacg
      acagcgcttc ggagatgggtg gtggagaagc accgggagtc ccgcggctgg
      gtggagaccc tgcggccgcg ctacacctac ttcaagggtgc ccacggagcg
      cgacctgggtc tactacgagg cctcgcccaa cttctgcgag cccaaccctg
25     agacgggctc cttcggcacg cgcgaccgca cctgcaacgt cagctcgcac
      ggcacgcacg gctgcgacct gctgtgctgc ggccgcggcc acaacgcgcg
      agcggagcgg cgccgggaga agtgccgctg cgtgtttcac tgggtgctg
      (SEQ ID NO:11)

```

Stem cells may be obtained from a a heterologous

30 donor animal such as a pig. The animal is euthanized and tissue removed using a sterile procedure. Brain areas of particular interest include any area from which progenitor cells can be obtained which will serve to restore function to a degenerated area of the host's

35 brain. These regions include areas of the CNS including the cerebral cortex, cerebellum, midbrain, brainstem, spinal cord and ventricular tissue, and areas of the peripheral nervous system (PNS) including the carotid body and the adrenal medulla. For example, cells may be

40 obtained from the basal ganglia, preferably the striatum which consists of the caudate and putamen, or various cell groups such as the globus pallidus, the subthalamic nucleus, or the substantia nigra pars compacta (which is found to be degenerated in Parkinson's Disease patients).

- 15 -

Human heterologous neural progenitor cells may be derived from fetal tissue obtained from elective abortion, or from a post-natal, juvenile or adult organ donor. Autologous neural tissue can be obtained by
5 biopsy, or from patients undergoing neurosurgery in which neural tissue is removed, in particular during epilepsy surgery, and more particularly during temporal lobectomies and hippocampalectomies.

Cells can be obtained from donor tissue by
10 dissociation of individual cells from the connecting extracellular matrix of the tissue. Dissociation can be obtained using any known procedure, including treatment with enzymes, e.g., trypsin or collagenase, or by using physical methods of dissociation such as with a blunt
15 instrument. Dissociation of fetal cells can be carried out in tissue culture medium, while a preferable medium for dissociation of juvenile and adult cells is artificial cerebral spinal fluid (aCSF). Regular aCSF contains 124 mM NaCl, 5 mM KCl, 1.3 mM MgCl₂, 2 mM CaCl₂,
20 26 mM NaHCO₃, and 10 mM D-glucose. Low Ca²⁺ aCSF contains the same ingredients except for MgCl₂ at a concentration of 3.2 mM and CaCl₂ at a concentration of 0.1 mM.

Dissociated cells can be placed into any culture medium capable of supporting cell growth, including MEM,
25 DMEM, RPMI, F-12. The medium may containin supplements which support cellular metabolism such as glutamine and other amino acids, vitamins, minerals and proteins such as transferrin. In some cases, the medium may contain bovine, equine, chicken or human serum. A preferable
30 medium for neural precursor cells is a mixture of DMEM and F-12. Conditions for culturing mimic physiological conditions, e.g., physiological pH, preferably between pH 6-8, more preferably close to pH 7, even more particularly about pH 7.4 at a temperature that is at or
35 close to physiological temperature.

- 16 -

Cells can be grown in suspension or on a fixed substrate, but proliferation of the precursor cells is preferably done in suspension to generate large numbers of cells by formation of "neurospheres" (see, for
5 example, Reynolds et al., 1992, Science 255:1070-1709; and PCT Publications WO93/01275, WO94/09119, WO94/10292, and WO94/16718). Cell suspensions in culture medium are supplemented with any growth factor which allows for the proliferation of precursor cells and seeded in any
10 receptacle capable of sustaining cells, preferably in culture flasks or roller bottles. Cells typically proliferate within 3-4 days in a 37°C incubator, and proliferation can be reinitiated at any time after that by dissociation of the cells and resuspension in fresh
15 medium containing growth factors.

In the absence of substrate, cells lift off the floor of the flask and continue to proliferate in suspension forming a hollow sphere of undifferentiated cells. After approximately 3-10 days *in vitro*, the
20 proliferating clusters (neurospheres) are fed every 2-7 days, and more particularly every 2-4 days by gentle centrifugation and resuspension in medium containing a Wnt polypeptide or a growth factor.

After 6-7 days *in vitro*, individual cells in the
25 neurospheres can be separated by physical dissociation of the neurospheres with a blunt instrument, more particularly by titrating the neurospheres with a pipette. Single cells from the dissociated neurospheres are suspended in culture medium containing growth
30 factors, and differentiation of the cells can be induced by plating (or resuspending) the cells in the presence of a Wnt agonist, and (optionally) any other factor capable of inducing and/or sustaining differentiation.

The tissue culture media is supplemented with a
35 Wnt polypeptide (either by adding a Wnt polypeptide to

- 17 -

the culture media or by adding feeder cells producing a Wnt polypeptide) to maintain a stem cell phenotype of the precursor cells and to promote proliferation of the cells. Other commercially available growth factors such as Fibroblast Growth Factor (FGF) or Epidermal Growth Factor (EGF) are added to the culture as mitogens.

Cells cultured in the presence of a Wnt polypeptide, e.g., a member of the Wnt-1 class of polypeptides, proliferate and maintain a stem cell phenotype. Differentiation of the cells can proceed upon the removal of the Wnt polypeptide and/or addition of a composition that promotes differentiation.

A naturally-occurring population of neural crest cells contains multipotent (i.e., uncommitted) neural crest cells and committed precursor cells. The role of Wnt proteins employed in the present method is to culture a population of neural precursor cells, e.g., a naturally-occurring population of neural crest cells, (1) to induce cell fate of an uncommitted precursor and thereby give rise to a committed precursor cell and (2) to maintain such cells in a stem cell state (e.g., to arrest the development of a committed precursor cell towards becoming a terminally-differentiated neuronal cell). For example, the present method can be used in vitro to induce and/or maintain the differentiation of neural crest cells into glial cells, schwann cells, chromaffin cells, cholinergic sympathetic or parasympathetic neurons, as well as peptidergic and serotonergic neurons. The Wnt protein can be used alone, or can be used in combination with other neurotrophic factors which act to more particularly enhance a particular differentiation fate of the neuronal precursor cell. In the later instance, an Wnt polypeptide might be viewed as ensuring that the treated cell has achieved a particular phenotypic state such that the cell is poised

- 18 -

along a certain developmental pathway so as to be properly induced upon contact with a secondary neurotrophic factor. Even relatively undifferentiated stem cells or primitive neuroblasts can be maintained in culture and caused to differentiate by treatment with Wnt agonists. Exemplary primitive cell cultures comprise cells harvested from the neural plate or neural tube of an embryo.

A population of neural precursor cells is characterized as having a stem cell phenotype when the level of proliferation of the cells in standard tissue culture media (e.g., MEM, DMEM, RPMI, F-12) in the presence of a Wnt polypeptide is at least 20% greater than the level of proliferation in the same tissue culture media without the Wnt polypeptide. Preferably, the level of cell proliferation is at least 50% greater in the presence of a Wnt polypeptide compared to the level of proliferation in the absence of a Wnt polypeptide. Proliferation is measured using known methods, e.g., incorporation of tritiated thymidine. Neural cells with a differentiated phenotype are characterized as non-proliferating and having the physical characteristics and cell markers of a mature terminally-differentiated neuron.

Primary stem cells may be immortalized by a variety of known techniques such as retrovirus-mediated transduction of an immortalizing gene, e.g., avian *myc* (*v-myc*).

Graft preparation

The therapeutic methods of the invention which utilize enriched populations of neural precursor cells may be used to treat neurodegenerative diseases and other types of diseases that result in depletion of neural cells. In addition to chronic depletion associated with progressive neurodegenerative diseases, neurons may be

- 19 -

killed as a consequence of infectious diseases, autoimmune diseases, and immunodeficiency diseases. Clinical outcome of treatment can be assessed by measuring as motor and cognitive capabilities of the patient, length of patient survival, quality of life.

Precursor cells cultured in the presence of a Wnt polypeptide as described above are washed and resuspended in a pharmaceutically acceptable excipient, e.g., a solution of 0.6% glucose-saline, are transplanted into brain tissue of a recipient mammal using known methods, e.g., those described by Gage et al., 1987, Ciba Found. Symp. 126:143-159. A small volume of a cell suspension is stereotactically injected into a desired region, e.g., the hippocampus, of a mammal. For example, approximately 10⁶ cells are infused into a desired location of the brain of the patient over 30 min.

Subsequent to transplantation, a Wnt polypeptide may be administered to the patient to induce further proliferation of stem cell *in vivo*. Wnt polypeptides can be administered in the form of a nerve prostheses for the repair of central and peripheral nerve damage. In particular, where a crushed or severed axon is intubulated by use of a prosthetic device, Wnt polypeptides can be added to the prosthetic device to increase the rate of growth and regeneration of the dendritic processes.

Alternatively, prior to transplantation, the cells may be exposed to a composition that induces differentiation Treatment of neurodegenerative disease

Neurodegenerative diseases include familial and sporadic amyotrophic lateral sclerosis (FALS and ALS, respectively), familial and sporadic Parkinson's disease, Huntington's disease, familial and sporadic Alzheimer's disease, olivopontocerebellar atrophy, multiple system atrophy, progressive supranuclear palsy, diffuse lewy

- 20 -

body disease, corticodentatonigral degeneration, progressive familial myoclonic epilepsy, strionigral degeneration, torsion dystonia, familial tremor, gilles de la tourette syndrome, and Hallervorden-Spatz disease.

5 Most of the diseases are typified by onset during the middle adult years and lead to rapid degeneration of specific subsets of neurons within the neural system, ultimately resulting in premature death. There is no known cure nor is there an effective therapy to slow the
10 progression for any of the listed diseases.

Parkinson's disease (paralysis agitans) is a common neurodegenerative disorder which appears in mid to late life. Familial and sporadic cases occur, although familial cases account for only 1-2 percent of the
15 observed cases. The neurological changes which cause this disease are somewhat variable and not fully understood. Patients frequently have nerve cell loss with reactive gliosis and Lewy bodies in the substantia nigra and locus coeruleus of the brain stem. Similar
20 changes are observed in the nucleus basalis of Meynert. Nigrostriatal dopaminergic neurons are most affected.

The disorder generally develops asymmetrically with tremors in one hand or leg and progresses into symmetrical loss of voluntary movement. Eventually, the
25 patient becomes incapacitated by rigidity and tremors. In the advanced stages the disease is frequently accompanied by dementia.

Diagnosis of both familial and sporadic cases of Parkinson's disease can only be made after the onset of
30 the disease. Anticholinergic compounds, propranolol, primidone and levodopa are frequently administered to modify neural transmissions and thereby suppress the symptoms of the disease, though there is no known therapy which halts or slows the underlying progression. The
35 therapeutic methods described herein may be administered

- 21 -

in conjunction with existing therapeutic approaches to neurodegenerative diseases.

The death of the dopaminergic neurons in the basal ganglia is an underlying defect of this progressive chronic disease as the basal ganglia are involved in the control of voluntary movements. Wnt-polypeptides and neural precursor cells cultured in the presence of Wnt polypeptides, e.g., Wnt-1, are useful in the treatment of Parkinson's disease and other disorders of midbrain dopamine circuitry. Transplantation of dopaminergic neural precursor cells is used to repopulate a patient's depleted population of dopaminergic neurons to treat or ameliorate the symptoms of Parkinson's disease.

Another neurodegenerative disease, Alzheimer's disease, can take two forms: disease exist: presenile dementia, in which the symptoms emerge during middle age, and senile dementia which occurs in the elderly. Both forms of the disease appear to have the same pathology. Diseases which affect learning and memory may be characterized by a depletion of hippocampal cells. Transplantation of hippocampal neural precursor cell is used to repopulate a patient's depleted population of hippocampal neurons to treat neurodegenerative diseases that affect learning and memory such as Alzheimer's disease.

Example 1: Wnt Signaling and Proliferation

Wnt signalling was found to regulate the expansion of dorsal neural precursors. Wnt-1 and Wnt-3a are coexpressed at the dorsal midline of the developing neural tube. Wnt-1 is involved in midbrain patterning, and Wnt-3a is involved in the formation of the paraxial mesoderm. The absence of a dorsal neural tube phenotype in animals with a mutation in either gene suggested that Wnt signalling is redundant. The data described below indicate that in the absence of both Wnt-1 and Wnt-3a,

- 22 -

there is a marked deficiency in neural crest derivatives, which originate from the dorsal neural tube, and a pronounced reduction in dorsolateral precursors within the neural tube itself.

5 Mice lacking both Wnt-1 and Wnt-3a signaling were generated. Mice which are heterozygous for null alleles of Wnt-1 and Wnt-3a were made using known methods (e.g., McMahon et al., 1990, Cell 62:1073-1085 and Takada et al., 1994, Genes Dev. 8:174-189). Compound heterozygotes
10 (on a predominantly 129/Sv background) were intercrossed to recover compound mutants. Genotypes were confirmed by genomic Southern hybridization and polymerase chain reaction (PCR). Whole mount immunostaining was carried out using antibodies specific for neurofilaments, CRABP-
15 1, and Lmx-1b. Skeletons from 18.5 d.p.c embryos were prepared and stained with alcian blue and alizarin red using known methods.

To evaluate cell proliferation and death, embryos were collected at 9.5 d.p.c (20-25 somite stage
20 development) after intraperitoneal injection of pregnant females with 50 μ g per body weight of 5-bromo-2'-deoxyuridine (BrdU). Mice were killed one hour later. Embryos were fixed overnight in 4% paraformaldehyde in phosphate-buffered saline (PBS) at 4°C. After
25 dehydration, wax embedding and sectioning at a thickness of 6 μ m, serial sections were dewaxed and either stained with haematoxylin and eosin, or assayed for BrdU incorporation for apoptotic death using a standard TUNEL procedure.

30 Compound homozygotes were recovered at the expected Mendelian frequency (51 compound homozygotes in 673 embryos. The frequency was close to the expected frequency of 1/16) between 9.0 and 10.5 days post coitum (d.p.c.). Due to the termination of caudal axial
35 development accompanying the loss of Wnt-3a activity,

- 23 -

relatively few of these embryos survived to 18.5 d.p.c. (3 compound homozygotes in 151 embryos).

To assess the development of the dorsal neural tube in compound mutants, neural crest derived structures were examined. Neural crest cells are among the first differentiated cell types to be formed by dorsal neural precursors. Neurofilament staining indicated that both neural crest derived cranial and spinal ganglia formation were unaltered in single mutants (either Wnt-1 or Wnt-3a mutants) which were either wild type or heterozygous for mutations in the other Wnt member. However, in double mutants, neurons derived from the proximal ganglion of cranial nerve IX (glossopharyngeal), which is formed by crest cells originating from rhombomere 6 within the hindbrain (r6), were absent. In contrast, the distal ganglion which is placodal in origin was present. In addition, there was a marked reduction in the proximal axons of cranial nerves V (trigeminal, r2 derived) and X (vagus, r7 derived). Similarly, in the trunk, there was a reduction in neurofilament staining in the cervical dorsal root ganglia. Further, cell counts indicated a 60% decrease in the cellular content of the dorsal root ganglia. Whole-mount *in situ* hybridization with probes specific for *Islet-1* and *cadherin-6*, markers of neuronal and glial neural crest derivatives, respectively, confirmed the reduction or absence of crest cells within the cranial ganglia and dorsal root ganglia. In contrast sympathetic ganglia, which express *c-ret*, were unaffected.

The reduction of neurogenic and gliogenic crest derivatives in the caudal head and rostral trunk regions indicates that fewer neural crest cells emerge in embryos lacking both Wnt-1 and Wnt-3a signaling. The issue of neural crest formation was evaluated by examining CRABP-1 immunoreactivity and AP-2 transcription. CRABP-1 is

- 24 -

normally present in the dorsal CNS at 9.0 d.p.c. as well as in migrating neural crest cells arising from r2, 4 and 6. AP-2 is first expressed at 8.5 d.p.c. in the dorsal neural plate, coincident with neural crest formation. By 5 9.5 d.p.c. cranial expression is absent in the neural tube but persists in migrating and maturing neural crest derivatives at cranial and spinal cord levels. Loss of function studies have demonstrated that AP-2 is essential for development of neural crest derived structures. A 10 clear decrease was observed in migrating CRABP-1 positive cells within the hindbrain, although CRABP-1 staining within the CNS appeared to be relatively normal. Similarly, examination of AP-2 expression revealed a reduction in both cranial and trunk neural crest. In 15 contrast to their wild type litter mates, double mutants also retained AP-2 expression within the dorsal CNS at 9.5 d.p.c. Thus, in the absence of Wnt-1 and Wnt-3a, there is both a reduction in neural crest cell formation and persistent expression of AP-2 at the dorsal midline. 20 To determine whether Wnt-signaling was required throughout the period of cranial crest formation, expression of TRP-2 was evaluated. TRP-2 is a marker of presumptive melanocytes which are dominant in late formed cranial crest derivatives. At 11.5 d.p.c., TRP-2 25 expression was virtually absent within presumptive melanocytes migrating within the hindbrain region of double mutants though a few TRP-2 cells remained at the dorsal midline. In view of the prolonged expression of AP-2 within the dorsal CNS, TRP-2 expressing cells may be 30 differentiating at a later stage, or they may be retained at the midline because Wnt-signaling promotes neural crest migration. Neither CRABP-1, TRP-2 or AP-2 expression was altered in the forebrain indicating that there is regional specificity in the requirement for 35 these Wnt-signals.

- 25 -

Much of the head skeleton is generated by cranial neural crest. Distinct skeletal elements are derived from neural crest cells which emerge from different regions of the brain. To determine whether the reduction
5 in neural crest formation in double mutants leads to alterations in the skeleton, 18.5 d.p.c. embryos were stained with alcian blue and alizarin red to examine cartilage and bone development. The stapes and the main body of the hyoid bone including the greater horn which
10 originate from crest cells derived from r3-5 and r6-7, respectively, were absent. Thyroid cartilage showed a consistent dysmorphology. The reduction in hindbrain crest formation was also reflected in the absence of specific skeletal derivatives. In contrast, despite the
15 early loss of forebrain, midbrain and rostral hindbrain in double mutants, the development of skeletal crest derivatives from these regions, as well as non-crest derived bones, was largely normal though there was some reduction in development of the squamosal, alisphenoid,
20 basisphenoid, presphenoid and otic capsule. These data indicate that, in the absence of Wnt-1/3a signaling, several neural crest cell fates form, but there is a dramatic reduction in neural crest derivatives in the hindbrain region and in the spinal cord.

25 Neural crest cell development, and other aspects of dorsal polarity within the developing CNS, are thought to be regulated by BMP signals produced initially by the dorsal ectoderm and subsequently by the dorsal CNS. BMP-7 expression was induced, as expected, in the roof plate
30 of double mutants. The data indicate that it was unlikely that defective neural crest development resulted from a secondary loss of BMP-signaling within the dorsal neural tube.

To determine whether Wnt-signaling directly
35 regulates dorso-ventral polarity within the CNS, the

- 26 -

distribution of a number of regionally expressed markers was examined. Whereas spinal cord levels appeared normal, the hindbrain displayed a striking phenotype. Expression of Wnt-3a, Wnt-1 and Lmx-1b was normal in the
5 roof plate. Thus, unlike other aspects of Wnt-signaling in the mammalian embryo, these Wnt-expressing cells did appear to require the Wnt-signals they produce. In contrast, expression of Math1 (which is activated at 9.5 d.p.c. in cells immediately adjacent to the roof plate)
10 and Pax-3 (which occupies most of the dorsal half of the CNS) were dramatically reduced in the double mutant hindbrain. Dbx expression at the dorsal-ventral interface and Pax-6 expression in the ventro-lateral CNS were normal.

15 The data indicate that in the hindbrain, Wnt-signaling does not appear to play a role directly in the primary patterning processes which lead to the establishment of distinct cell fates in appropriate positions along the dorsoventral axis. Rather, it
20 appears to play an essential role in the subsequent expansion of dorso-lateral neural progenitors. In support of a potential role in neural proliferation, transgenic analysis demonstrated that Wnt-1 can act as a potent mitogen when ectopically expressed within the
25 dorsal CNS.

 In normal development there is a ventral to dorsal progression in the formation of different neural crest derivatives. In the double mutants, the most severely affected crest derivatives were more proximal (dorsally
30 located) structures. The stapes was absent from the second branchial arch while the lesser horn of the hyoid was unaltered, and in the trunk, dorsal root ganglia were markedly reduced while the sympathetic ganglia appeared normal. If the signals governing commitment to neural
35 crest cell fates were unperturbed in the double mutant,

- 27 -

but renewal of a multipotential dorsal neural progenitor pool required Wnt-signals, the expected result would be a loss of later forming neural crest derivatives in Wnt-1/-3a mutants, as precursors within the neural tube became
5 limiting.

Cell proliferation and cell death in hindbrain tissue sections (9.5 d.p.c; 20-25 somites) were analyzed using BrdU incorporation and TUNEL staining, respectively.

10 Dorsal neural precursors were reduced, but no discernible change was detected in either proliferation or cell death within remaining dorsal regions of Wnt-1 and Wnt-3a mutants. As these two Wnts are first coexpressed at the otic level when the first neural crest cells appear (at
15 about 8.5 d.p.c; 8-10 somites), it is likely that the main reduction in dorsolateral neural precursors occurs between 8.5 and 9.5 d.p.c.

These data indicate that Wnt signalling regulates dorsoventral patterning in the mammalian CNS through the
20 control of cell proliferation.

Example 2: Wnt-3A Signaling in Neuronal Differentiation

Wnt-3a expression in the mouse begins in the primitive streak region of the late egg cylinder at 7.5 d.p.c. and is maintained in the tail bud until tail
25 formation is complete. To determine which cell types in the primitive streak region express Wnt-3a, the expression of Wnt-3a transcripts was examined in wild type embryos at the 7 somite stage. Expression was detected in the ectoderm layer in the primitive streak
30 region but was absent from the node. Expression was further restricted for ventrally located cells in the anterior streak region. In contrast, in the posterior streak, most cells in the ectoderm layer expressed Wnt-3a. Wnt-3a expression was not observed in migrating
35 mesodermal cells at either anterior or posterior

- 28 -

positions. These data indicate that Wnt-3a expression is localized to the primitive ectoderm prior to the physical segregation of the paraxial mesoderm and is downregulated as cells ingress through the primitive streak.

5 The phenotype of Wnt-3a homozygous mutant embryos was analyzed at early somite stages. At the 5 somite stage, no obvious differences in morphology between wild type and Wnt-3a mutant embryos were detected. However, by the 7 somite stage, differences in the shape of the
10 primitive streak region were apparent. In Wnt-3a mutants, the width of the primitive streak region is narrower than in the wild type embryos and this phenotype becomes more pronounced by the 10 somite stage.

 To further investigate the abnormal morphology of
15 mutant embryo, histological analysis of the sections was carried out. In wild type embryos at the 7 somite stage, migrating presomitic mesodermal cells were observed under the primitive ectoderm layer in the primitive streak region. However, in Wnt-3a mutant embryos at the same
20 stage, no migrating presomitic mesoderm cells were observed; in contrast, the shape and movement of cells ingressed through the primitive streak are quite different from those in normal embryos. In the anterior streak region of the mutant embryos, the ingressing cells
25 were round in appearance, quite distinct from the usual stellate mesenchymal morphology of the ingressing mesoderm. Furthermore, these cells contacted each other and formed an ectopic tubular structure under the primitive streak at more posterior level. This tubular
30 structure was not observed anterior to the streak where somites are present. Thus, in Wnt-3a mutant embryos, the absence of somite precursors appears to be correlated with the appearance of an ectopic tubular structure arising in the primitive streak region.

- 29 -

To identify the molecular characteristics of the ectopic tubular structure in Wnt-3a mutant embryos, *in situ* hybridization and whole mount immunostaining and the expression of a variety of molecular markers detected.

5 MF-1, encodes a forkhead domain containing protein, which is normally expressed in somites, presomitic mesoderm, and lateral mesoderm at 9.5 d.p.c. In Wnt-3a mutant embryos at this stage, no obvious MF-I expression was observed in the position where the ectopic
10 tube was formed posterior to the forelimb level. A transverse section of the stained embryo at this axial level clearly indicated that no MF-1 transcripts were localized in the ectopic tube. Similarly another paraxial mesoderm marker, Mox-1, was not expressed in the
15 ectopic tube in Wnt-3a mutants at either 8.5 or 9.5 d.p.c. The data indicate that the ectopic tube does not have the molecular and morphological characteristics of paraxial mesoderm.

Mash-I is normally expressed in central nervous
20 system and peripheral nervous system precursors at 9.5 d.p.c. but not in the mesoderm. In Wnt-3a mutant embryos at the same stage, *Mash-1* expression was detected not only in these region but also in the region ventral to the original neural tube posterior to the forelimb level.
25 A transverse section of Wnt-3a mutants at the axial level, where abnormal Mash-7 expression was observed, indicated that the ventral expression of Mash-I was localized in the ectopic tube. A second neural marker, HES-5, which is normally expressed in CNS, was also
30 expressed in the ectopic tube in Wnt-3a mutants at 9.5 d.p.c. To explore further whether neurons differentiate in the ectopic tube, Wnt-3a mutant embryos at 10.5 d.p.c. were immunostained with antineurofilament antibody, 2H3. Neurofilament expressing cells were present in both the
35 dorsal neural tube and the ectopic ventral tube.

- 30 -

The ectopic tube also exhibited polarity typical of CNS tissue. For example, Sonic hedgehog (Shh) is normally expressed in the floor plate of the neural tube. In 9.5 d.p.c. Wnt-3a mutant embryos, the notochord was present under the ventral ectopic tubular structure but not under the original neural tube at the axial level just posterior to the forelimbs while no notochord was absorbed at more posterior levels. Shh was expressed in ventrally in the ectopic tube where it contacts the notochord, suggesting, that the ectopic tube forms a floor plate in response to a Shh signaling by the notochord. The ectopic neural tube also exhibits dorsal polarity typical of the CNS such as the expression of the dorsal midline marker, Wnt-1 and increased levels of Pax-3 expression, where the tube contacts the surface ectoderm. In addition, expression of a ventral CNS marker, Pax-6, was suppressed where the ectopic tube contacts the surface ectoderm. Taken together, the data indicate that the ectopic tubular structure in the mutants has the molecular and cellular characteristics of an ectopic neural tube and consequently the loss of Wnt-3a signaling results in the formation of CNS precursors at the expense of paraxial mesoderm.

The phenotype of Wnt-3a knock out mutant embryos at 9.5 d.p.c. indicated that Wnt-3a is essential for formation of somitic mesoderm caudal to first 7-9 somites. In the absence of somite formation, an ectopic tubular structure which displays both cellular and molecular characteristics of presumptive CNS tissue is formed. Several lines of evidences suggest that the neural tube was formed ectopically. First, transverse sections of Wnt-3a mutant embryos at an early somite stage indicated that cells delaminating from and ingressing through the primitive streak form an epithelial cell layer that contribute to an ectopic tube

- 31 -

under the primitive ectoderm in the primitive streak region. Second, the notochord contacts the ventral but not the dorsal neural tube, suggesting that the ventral (ectopic) neural tube had already formed at the time when
5 the notochord was laid down. Third, by the analysis of serial transverse sections of several 8.5 and 9.5 d.p.c. Wnt-3a mutant embryos, it is apparent that the ectopic neural tube is not continuous with the original dorsal neural tube suggesting an independent origin.

10 The appearance of the ectopic neural tube correlates with the disappearance of the paraxial mesoderm precursors in Wnt-3a mutant embryos. This correlation suggests that the absence of Wnt-3a signaling in the primitive ectoderm of the primitive streak may
15 lead to presumptive somitic mesoderm cells to adopting, neural cell fate. That is, a neural fate may be a "default" state for cells which normally give rise to both mesodermal and neural derivatives.

 The results described herein indicate that in the
20 normal primitive ectoderm, where Wnt-3a is expressed, undifferentiated cells can differentiate into both neural and somitic mesoderm cell lineages. At early somite stages, cells in the anterior primitive streak generate mostly somitic mesoderm, while cells in the posterior
25 streak gives rise to mostly lateral mesoderm. In contrast, primitive ectoderm adjacent to the anterior primitive streak contributes mainly to somitic mesoderm and neuroectoderm, suggesting that these two cell types might arise from the same cell population. The data
30 indicate that Wnt-3a signaling regulates cell fate specification between somitic mesoderm and neural lineages in the normal mouse embryo.

 Although Wnt-3a is expressed in the anterior streak in regions which gives rise to somitic mesoderm,
35 it is also expressed in more posterior regions which

- 32 -

generate lateral and ventral mesoderm. Thus, expression is not restricted to paraxial mesoderm precursors. Wnt-3a may establish a competence to respond to a paraxial mesoderm inducing signal, rather than itself directly inducing paraxial mesodermal cell fates. This competence may be broadly distributed within the streak.

Example 3: Wnt-1 signaling and mid-brain development

Expression of En-1 in the developing midbrain of Wnt-1 null embryos is sufficient to rescue midbrain and interior hindbrain development. In the mouse, Wnt-1 and Engrailed-1 (En-1) are first expressed in the presumptive midbrain, from 8.0 days post coitum (d.p.c.) and continue to be expressed, together with En-2, in overlapping patterns during midbrain development. In Wnt-1^{-/-} (Wnt-1-null) embryos, En-1 and En-2 expression is initiated normally, but subsequently both domains of En expression are lost, which is concomitant with a failure of midbrain and anterior hindbrain development.

En-1 was expressed from the transgene WEXPZ-En-1 in a pattern similar to that of endogenous Wnt-1 gene. To assess whether En-1 was able to rescue the Wnt-1-null phenotype, embryos from matings of Wnt-1^{+/-}, WEXPZ-En-1⁺ males with Wnt-1^{+/-} females were collected at 14.5 d.p.c., when the Wnt-1^{-/-} phenotype can easily be scored morphologically. The genotype was subsequently determined by southern blotting. Wnt-1^{+/+} and Wnt-1^{+/-} embryos with or without WEXPZ-En-1 appeared to be wild-type (n = 112) whereas all Wnt-1^{-/-} embryos (n = 12) displayed the Wnt-1^{-/-} phenotype. In Wnt-1^{-/-}, WEXPZ-En-1⁺ embryos, 7 out of 17 appeared superficially wildtype, 8 out of 17 were partially rescued and only 2 out of 17 were similar to Wnt-1^{-/-} embryos.

To characterize brain development in greater detail, a minimum of four embryos from each category were

- 33 -

sectioned for histological analysis. All Wnt-1^{-/-} embryos lacked the midbrain and cerebellum. In contrast, in Wnt-1^{-/-}, WEXPZ-En-1⁺ embryos that were scored as wild-type, the midbrain and cerebellum appeared similar to those of
5 wild-type embryos. In partially rescued embryos, only the posterior midbrain and a slightly reduced cerebellum were apparent. The absence of rescue in some cases, and partial rescue in others, may reflect influences of the genetic background or variations in the levels of En-1
10 expressed from the transgene.

To characterize the development of the midbrain in Wnt-1^{-/-}, WEXPZ-En-1⁺ embryos further, the expression of several genes normally transcribed in this region was examined at 10.5 d.p.c. Pax-5 is expressed in a broad
15 domain at the midbrain-hindbrain junction, but this domain is missing in Wnt-1^{-/-} embryos. In Wnt-1^{-/-}, WEXPZ-En-1⁺ embryos, Pax-5 expression was detected in a pattern similar to that of the wild-type embryos. Wnt-1 and Fgf-8 are normally expressed in adjacent rings of cells just
20 anterior and posterior to the midbrain-hindbrain junction, respectively. Fgf8 signalling is involved in midbrain development. In Wnt-1^{-/-} embryos, both rings of expressing cells are absent. In contrast, both Wnt-1 and Fgf-8 were expressed in sharp rings of cells in Wnt-1^{-/-},
25 WEXPZ-En-1⁺ embryos despite the fact that no morphologically obvious midbrain-hindbrain junction was apparent. These results indicate that Wnt-1 signaling at this later stage may not play a direct role in regulating Fgf-8 expression in adjacent cells. En gene expression
30 was also restored in the mid-hindbrain region of Wnt-1^{-/-}, WEXPZ-En-1⁺ embryos outside the area where the transgene is expressed.

In all the rescued embryos, the expression domains of Pax-5, Fgf-8, En, and, in a few cases, Wnt-1 were

- 34 -

slightly reduced relative to wild-type littermates (18 out

41 Wnt-1^{-/-}, WEXPZ-En-1⁺ embryos expressed one of the markers examined, of these at least half were

5 substantially rescued). One likely explanation is that rescued embryos have a smaller population of midbrain cells than wild-type siblings because when Wnt-1 and En-1 expression is initiated, Wnt-1 mRNA transcription is patchy, whereas En genes are expressed more uniformly in
10 presumptive midbrain cells. Thus, in Wnt-1^{-/-}, WEXPZ-En-1⁺ embryos, where En-1 is not uniformly expressed in all presumptive midbrain cells, only those cells that express En-1 at this early stage may contribute to midbrain development. As En-1 expression in the midbrain restores
15 Fgf-8, Pax-5 and En expression in the anterior hindbrain, and subsequently cerebellum development in Wnt-1^{-/-} embryos, the data suggest that a midbrain-derived signal other than Wnt-1 is necessary for anterior hindbrain development.

20 To assess whether expression of En-1 was sufficient to rescue the viability of Wnt-1^{-/-} mice (pups are born but die within 24 h) pups were genotyped at 10 days post partum (n = 68). No live Wnt-1^{-/-}, WEXPZ-En-1⁺ mice were obtained indicating that En-1 was
25 insufficient to rescue the Wnt-1-null phenotype completely. Further analysis indicated that between 14.5 and 18.5 d.p.c., brains of Wnt-1^{-/-}, WEXPZ-En-1⁺ embryos deteriorate, indicating that there may be additional functions of Wnt-1 signaling that cannot be replaced by
30 En-1. This conclusion is supported by analysis of two cranial motor nerves, III (oculomotor) and IV (trochlear), which normally develop adjacent to Wnt-1-expressing cells in the ventral midbrain. Each of these fail to develop in Wnt-1^{-/-} embryos. Similarly, neither
35 nerve forms in Wnt-1^{-/-}, WEXPZ-En-1⁺ embryos which have

- 35 -

global restoration of midbrain development. In contrast, a second ventral population, tyrosine-hydroxylase-expressing neurons (catecholaminergic neurons) of the substantia nigra, are rescued in Wnt-1^{-/-}, WEXPZ-En-1⁺ embryos.

These data demonstrate that, in the absence of a Wnt-1 signal, expression of En-1 from the Wnt-1 enhancer is sufficient to substantially rescue early midbrain and anterior hindbrain development, and suggest that a major role of Wnt-1 signalling in the mammalian brain is to maintain En expression.

Other embodiments are within the following claims.

- 36 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: President and Fellows of Harvard College
- (ii) TITLE OF INVENTION: INDUCTION OF NEURONAL REGENERATION
- (iii) NUMBER OF SEQUENCES: 11
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Fish & Richardson P.C.
 - (B) STREET: 225 Franklin Street
 - (C) CITY: Boston
 - (D) STATE: MA
 - (E) COUNTRY: USA
 - (F) ZIP: 02110-2804
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Diskette
 - (B) COMPUTER: IBM Compatible
 - (C) OPERATING SYSTEM: Windows 95
 - (D) SOFTWARE: FastSEQ for Windows Version 2.0b
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER: PCT/US98/-----
 - (B) FILING DATE: 30-APR-1998
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Freeman, John W.
 - (B) REGISTRATION NUMBER: 29,066
 - (C) REFERENCE/DOCKET NUMBER: 00246/222WO1
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 617/542-5070
 - (B) TELEFAX: 617/542-8906
 - (C) TELEX: 200154

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 370 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

```

Met Gly Leu Trp Ala Leu Leu Pro Gly Trp Val Ser Ala Thr Leu Leu
 1           5           10           15
Leu Ala Leu Ala Ala Leu Pro Ala Ala Leu Ala Ala Asn Ser Ser Gly
           20           25           30
Arg Trp Trp Gly Ile Val Asn Val Ala Ser Ser Thr Asn Leu Leu Thr
           35           40           45
Asp Ser Lys Ser Leu Gln Leu Val Leu Glu Pro Ser Leu Gln Leu Leu
           50           55           60
Ser Arg Lys Gln Arg Arg Leu Ile Arg Gln Asn Pro Gly Ile Leu His

```

- 37 -

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 65 | | | | | 70 | | | | | 75 | | | | | 80 |
| Ser | Val | Ser | Gly | Gly | Leu | Gln | Ser | Ala | Val | Arg | Glu | Cys | Lys | Trp | Gln |
| | | | | 85 | | | | | 90 | | | | | 95 | |
| Phe | Arg | Asn | Arg | Arg | Trp | Asn | Cys | Pro | Thr | Ala | Pro | Gly | Pro | His | Leu |
| | | | 100 | | | | | 105 | | | | | 110 | | |
| Phe | Gly | Lys | Ile | Val | Asn | Arg | Gly | Cys | Arg | Glu | Thr | Ala | Phe | Ile | Phe |
| | | 115 | | | | | 120 | | | | | 125 | | | |
| Ala | Ile | Thr | Ser | Ala | Gly | Val | Thr | His | Ser | Val | Ala | Arg | Ser | Cys | Ser |
| | 130 | | | | | 135 | | | | | 140 | | | | |
| Glu | Gly | Ser | Ile | Glu | Ser | Cys | Thr | Cys | Asp | Tyr | Arg | Arg | Arg | Gly | Pro |
| 145 | | | | | 150 | | | | | 155 | | | | | 160 |
| Gly | Gly | Pro | Asp | Trp | His | Trp | Gly | Gly | Cys | Ser | Asp | Asn | Ile | Asp | Phe |
| | | | 165 | | | | | | 170 | | | | 175 | | |
| Gly | Arg | Leu | Phe | Gly | Arg | Glu | Phe | Val | Asp | Ser | Gly | Glu | Lys | Gly | Arg |
| | | 180 | | | | | 185 | | | | | 190 | | | |
| Asp | Leu | Arg | Phe | Leu | Met | Asn | Leu | His | Asn | Asn | Glu | Ala | Gly | Arg | Thr |
| | 195 | | | | | 200 | | | | | 205 | | | | |
| Thr | Val | Phe | Ser | Glu | Met | Arg | Gln | Glu | Cys | Lys | Cys | His | Gly | Met | Ser |
| | 210 | | | | | 215 | | | | | 220 | | | | |
| Gly | Ser | Cys | Thr | Val | Arg | Thr | Cys | Trp | Met | Arg | Leu | Pro | Thr | Leu | Arg |
| 225 | | | | | 230 | | | | | 235 | | | | | 240 |
| Ala | Val | Gly | Asp | Val | Leu | Arg | Asp | Arg | Phe | Asp | Gly | Ala | Ser | Arg | Val |
| | | | 245 | | | | | | 250 | | | | 255 | | |
| Leu | Tyr | Gly | Asn | Arg | Gly | Ser | Asn | Arg | Ala | Ser | Arg | Ala | Glu | Leu | Leu |
| | | | 260 | | | | 265 | | | | | 270 | | | |
| Arg | Leu | Glu | Pro | Glu | Asp | Pro | Ala | His | Lys | Pro | Pro | Ser | Pro | His | Asp |
| | 275 | | | | | 280 | | | | | | 285 | | | |
| Leu | Val | Tyr | Phe | Glu | Lys | Ser | Pro | Asn | Phe | Cys | Thr | Tyr | Ser | Gly | Arg |
| | 290 | | | | | 295 | | | | | 300 | | | | |
| Leu | Gly | Thr | Ala | Gly | Thr | Ala | Gly | Arg | Ala | Cys | Asn | Ser | Ser | Ser | Pro |
| 305 | | | | | 310 | | | | | 315 | | | | | 320 |
| Ala | Leu | Asp | Gly | Cys | Glu | Leu | Leu | Cys | Cys | Gly | Arg | Gly | His | Arg | Thr |
| | | | 325 | | | | | | 330 | | | | 335 | | |
| Arg | Thr | Gln | Arg | Val | Thr | Glu | Arg | Cys | Asn | Cys | Thr | Phe | His | Trp | Cys |
| | | 340 | | | | | 345 | | | | | 350 | | | |
| Cys | His | Val | Ser | Cys | Arg | Asn | Cys | Thr | His | Thr | Arg | Val | Leu | His | Glu |
| | | 355 | | | | 360 | | | | | | 365 | | | |
| Cys | Leu | | | | | | | | | | | | | | |
| | 370 | | | | | | | | | | | | | | |

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 360 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Met | Asn | Ala | Pro | Leu | Gly | Gly | Ile | Trp | Leu | Trp | Leu | Pro | Leu | Leu | Leu |
| 1 | | | | 5 | | | | | 10 | | | | | 15 | |
| Thr | Trp | Leu | Thr | Pro | Glu | Val | Asn | Ser | Ser | Trp | Trp | Tyr | Met | Arg | Ala |
| | | 20 | | | | | 25 | | | | | 30 | | | |
| Thr | Gly | Gly | Ser | Ser | Arg | Val | Met | Cys | Asp | Asn | Val | Pro | Gly | Leu | Val |
| | | 35 | | | | 40 | | | | | 45 | | | | |
| Ser | Ser | Gln | Arg | Gln | Leu | Cys | His | Arg | His | Pro | Asp | Val | Met | Arg | Ala |
| | 50 | | | | 55 | | | | | 60 | | | | | |
| Ile | Ser | Gln | Gly | Val | Ala | Glu | Trp | Thr | Ala | Glu | Cys | Gln | His | Gln | Phe |
| 65 | | | | 70 | | | | | 75 | | | | | 80 | |
| Arg | Gln | His | Arg | Trp | Asn | Cys | Asn | Thr | Leu | Asp | Arg | Asp | His | Ser | Leu |

| | | | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|--|--|
| | | | | 85 | | | | | 90 | | | | | 95 | | | |
| Phe | Gly | Arg | Val | Leu | Leu | Arg | Ser | Ser | Arg | Glu | Ser | Ala | Phe | Val | Tyr | | |
| | | | 100 | | | | | 105 | | | | | 110 | | | | |
| Ala | Ile | Ser | Ser | Ala | Gly | Val | Val | Phe | Ala | Ile | Thr | Arg | Ala | Cys | Ser | | |
| | | 115 | | | | | 120 | | | | | 125 | | | | | |
| Gln | Gly | Glu | Val | Lys | Ser | Cys | Ser | Cys | Asp | Pro | Lys | Lys | Met | Gly | Ser | | |
| | 130 | | | | | 135 | | | | | 140 | | | | | | |
| Ala | Lys | Asp | Ser | Lys | Gly | Ile | Phe | Asp | Trp | Gly | Gly | Cys | Ser | Asp | Asn | | |
| 145 | | | | | 150 | | | | | 155 | | | | | 160 | | |
| Ile | Asp | Tyr | Gly | Ile | Lys | Phe | Ala | Arg | Ala | Phe | Val | Asp | Ala | Lys | Glu | | |
| | | | 165 | | | | | | 170 | | | | | 175 | | | |
| Arg | Lys | Gly | Lys | Asp | Ala | Arg | Ala | Leu | Met | Asn | Leu | His | Asn | Asn | Arg | | |
| | | | 180 | | | | | 185 | | | | | 190 | | | | |
| Ala | Gly | Arg | Lys | Ala | Val | Lys | Arg | Phe | Leu | Lys | Gln | Glu | Cys | Lys | Cys | | |
| | | 195 | | | | | 200 | | | | | 205 | | | | | |
| His | Gly | Val | Ser | Gly | Ser | Cys | Thr | Leu | Arg | Thr | Cys | Trp | Leu | Ala | Met | | |
| | 210 | | | | | 215 | | | | | 220 | | | | | | |
| Ala | Asp | Phe | Arg | Lys | Thr | Gly | Asp | Tyr | Leu | Trp | Arg | Lys | Tyr | Asn | Gly | | |
| 225 | | | | | 230 | | | | | 235 | | | | | 240 | | |
| Ala | Ile | Gln | Val | Val | Met | Asn | Gln | Asp | Gly | Thr | Gly | Phe | Thr | Val | Ala | | |
| | | | 245 | | | | | | 250 | | | | | 255 | | | |
| Asn | Glu | Arg | Phe | Lys | Lys | Pro | Thr | Lys | Asn | Asp | Leu | Val | Tyr | Phe | Glu | | |
| | | | 260 | | | | | 265 | | | | | 270 | | | | |
| Asn | Ser | Pro | Asp | Tyr | Cys | Ile | Arg | Asp | Arg | Glu | Ala | Gly | Ser | Leu | Gly | | |
| | | 275 | | | | | 280 | | | | | 285 | | | | | |
| Thr | Ala | Gly | Arg | Val | Cys | Asn | Leu | Thr | Ser | Arg | Gly | Met | Asp | Ser | Cys | | |
| | 290 | | | | | 295 | | | | | 300 | | | | | | |
| Glu | Val | Met | Cys | Cys | Gly | Arg | Gly | Tyr | Asp | Thr | Ser | His | Val | Thr | Arg | | |
| 305 | | | | | 310 | | | | | 315 | | | | | 320 | | |
| Met | Thr | Lys | Cys | Gly | Cys | Lys | Phe | His | Trp | Cys | Cys | Ala | Val | Arg | Cys | | |
| | | | 325 | | | | | | 330 | | | | | 335 | | | |
| Gln | Asp | Cys | Leu | Glu | Ala | Leu | Asp | Val | His | Thr | Cys | Lys | Ala | Pro | Lys | | |
| | | | 340 | | | | | 345 | | | | | 350 | | | | |
| Asn | Ala | Asp | Trp | Thr | Thr | Ala | Thr | | | | | | | | | | |
| | | 355 | | | | | 360 | | | | | | | | | | |

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 352 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

| | | | | | | | | | | | | | | | |
|-----------|-----|-----------|------------|-----------|-----------|-----------|-----------|------------|-----------|-----------|-----|-----------|------------|-----------|-----------|
| Met 1 | Ala | Pro | Leu | Gly 5 | Tyr | Leu | Leu | Val | Leu 10 | Cys | Ser | Leu | Lys | Gln 15 | Ala |
| Leu | Gly | Ser | Tyr 20 | Pro | Ile | Trp | Trp | Ser 25 | Leu | Ala | Val | Gly | Pro 30 | Gln | Tyr |
| Ser | Ser | Leu 35 | Ser | Thr | Gln | Pro | Ile 40 | Leu | Cys | Ala | Ser | Ile 45 | Pro | Gly | Leu |
| Val 50 | Pro | Lys | Gln | Leu | Arg | Phe 55 | Cys | Arg | Asn | Tyr 60 | Val | Glu | Ile | Met | Pro |
| Ser 65 | Val | Ala | Glu | Gly | Val 70 | Lys | Ala | Gly | Ile | Gln 75 | Glu | Cys | Gln | His | Gln 80 |
| Phe | Arg | Gly | Arg | Arg 85 | Trp | Asn | Cys | Thr | Thr 90 | Val | Ser | Asn | Ser | Leu 95 | Ala |
| Ile | Phe | Gly | Pro 100 | Val | Leu | Asp | Lys | Ala 105 | Thr | Arg | Glu | Ser | Ala 110 | Phe | Val |
| His | Ala | Ile | Ala | Ser | Ala | Gly | Val | Ala | Phe | Ala | Val | Thr | Arg | Ser | Cys |

- 39 -

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| | | 115 | | | | | 120 | | | | | 125 | | | |
| Ala | Glu | Gly | Ser | Ala | Ala | Ile | Cys | Gly | Cys | Ser | Ser | Arg | Leu | Gln | Gly |
| | 130 | | | | | 135 | | | | | 140 | | | | |
| Ser | Pro | Gly | Glu | Gly | Trp | Lys | Trp | Gly | Gly | Cys | Ser | Glu | Asp | Ile | Glu |
| 145 | | | | | 150 | | | | | 155 | | | | | 160 |
| Phe | Gly | Gly | Met | Val | Ser | Arg | Glu | Phe | Ala | Asp | Ala | Arg | Glu | Asn | Arg |
| | | | 165 | | | | | | 170 | | | | | 175 | |
| Pro | Asp | Ala | Arg | Ser | Ala | Met | Asn | Arg | His | Asn | Asn | Glu | Ala | Gly | Arg |
| | | | 180 | | | | | | 185 | | | | 190 | | |
| Gln | Ala | Ile | Ala | Ser | His | Met | His | Leu | Lys | Cys | Lys | Cys | His | Gly | Leu |
| | | 195 | | | | | 200 | | | | | 205 | | | |
| Ser | Gly | Ser | Cys | Glu | Val | Lys | Thr | Cys | Trp | Trp | Ser | Gln | Pro | Asp | Phe |
| | 210 | | | | | 215 | | | | | 220 | | | | |
| Arg | Thr | Ile | Gly | Asp | Phe | Leu | Lys | Asp | Lys | Tyr | Asp | Ser | Ala | Ser | Glu |
| 225 | | | | | 230 | | | | | 235 | | | | | 240 |
| Met | Val | Val | Glu | Lys | His | Arg | Glu | Ser | Arg | Gly | Trp | Val | Glu | Thr | Leu |
| | | | | 245 | | | | | 250 | | | | | 255 | |
| Arg | Pro | Arg | Tyr | Thr | Tyr | Phe | Lys | Val | Pro | Thr | Glu | Arg | Asp | Leu | Val |
| | | | 260 | | | | | 265 | | | | | 270 | | |
| Tyr | Tyr | Glu | Ala | Ser | Pro | Asn | Phe | Cys | Glu | Pro | Asn | Pro | Glu | Thr | Gly |
| | | 275 | | | | | 280 | | | | | 285 | | | |
| Ser | Phe | Gly | Thr | Arg | Asp | Arg | Thr | Cys | Asn | Val | Ser | Ser | His | Gly | Ile |
| | 290 | | | | | 295 | | | | | 300 | | | | |
| Asp | Gly | Cys | Asp | Leu | Leu | Cys | Cys | Gly | Arg | Gly | His | Asn | Ala | Arg | Thr |
| 305 | | | | | 310 | | | | | 315 | | | | | 320 |
| Glu | Arg | Arg | Arg | Glu | Lys | Cys | His | Cys | Val | Phe | His | Trp | Cys | Cys | Tyr |
| | | | | 325 | | | | | 330 | | | | | 335 | |
| Val | Ser | Cys | Gln | Glu | Cys | Thr | Arg | Val | Tyr | Asp | Val | His | Thr | Cys | Lys |
| | | | 340 | | | | | 345 | | | | | 350 | | |

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 349 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

| | | | | | | | | | | | | | | | |
|------------|------------|------------|------------|-----------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|-----|
| Met 1 | Asn | Arg | Lys | Ala 5 | Leu | Arg | Cys | Leu | Gly 10 | His | Leu | Phe | Leu | Ser 15 | Leu |
| Gly | Met | Val | Cys 20 | Leu | Arg | Ile | Gly | Gly 25 | Phe | Ser | Ser | Val | Val 30 | Ala | Leu |
| Gly | Ala | Thr 35 | Ile | Ile | Cys | Asn | Lys 40 | Ile | Pro | Gly | Leu | Ala 45 | Pro | Arg | Gln |
| Arg | Ala 50 | Ile | Cys | Gln | Ser | Arg 55 | Pro | Asp | Ala | Ile | Ile 60 | Val | Ile | Gly | Glu |
| Gly 65 | Ser | Gln | Met | Gly 70 | Leu | Asp | Glu | Cys | Gln | Phe 75 | Gln | Phe | Arg | Asn 80 | Gly |
| Arg | Trp | Asn | Cys 85 | Ser | Ala | Leu | Gly | Glu | Arg 90 | Thr | Val | Phe | Gly | Lys 95 | Glu |
| Leu | Lys | Val 100 | Gly | Ser | Arg | Asp | Gly | Ala 105 | Phe | Thr | Tyr | Ala | Ile 110 | Ile | Ala |
| Ala | Gly 115 | Val | Ala | His | Ala | Ile | Thr 120 | Ala | Ala | Cys | Thr | His 125 | Gly | Asn | Leu |
| Ser | Asp 130 | Cys | Gly | Cys | Asp | Lys 135 | Glu | Lys | Gln | Gly | Gln 140 | Tyr | His | Arg | Asp |
| Glu 145 | Gly | Trp | Lys | Trp | Gly 150 | Gly | Cys | Ser | Ala | Asp 155 | Ile | Arg | Tyr | Gly | Ile |
| Gly | Phe | Ala | Lys 165 | Val | Phe | Val | Asp | Ala | Arg 170 | Glu | Ile | Lys | Gln | Asn 175 | Ala |

- 40 -

```

Arg Thr Leu Met Asn Leu His Asn Asn Glu Ala Gly Arg Lys Ile Leu
      180      185      190
Glu Glu Asn Met Lys Leu Glu Cys Lys Cys His Gly Val Ser Gly Ser
      195      200      205
Cys Thr Thr Lys Thr Cys Trp Thr Thr Leu Pro Gln Phe Arg Glu Leu
      210      215      220
Gly Tyr Val Leu Lys Asp Lys Tyr Asn Glu Ala Val His Val Glu Pro
      225      230      235      240
Val Arg Ala Ser Arg Asn Lys Arg Pro Thr Phe Leu Lys Ile Lys Lys
      245      250      255
Pro Leu Ser Tyr Arg Lys Pro Met Asp Thr Asp Leu Val Tyr Ile Glu
      260      265      270
Lys Ser Pro Asn Tyr Cys Glu Glu Asp Pro Val Thr Gly Ser Val Gly
      275      280      285
Thr Gln Gly Arg Ala Cys Asn Lys Thr Ala Pro Gln Ala Ser Gly Cys
      290      295      300
Asp Leu Met Cys Cys Gly Arg Gly Tyr Asn Thr His Gln Tyr Ala Arg
      305      310      315      320
Val Trp Gln Cys Asn Cys Lys Phe His Trp Cys Cys Tyr Val Lys Cys
      325      330      335
Asn Thr Cys Ser Glu Arg Thr Glu Met Tyr Thr Cys Lys
      340      345

```

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 124 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

```

Gly Val Ser Gly Ser Cys Thr Thr Lys Thr Cys Trp Thr Thr Leu Pro
  1      5      10      15
Lys Phe Arg Glu Val Gly His Leu Leu Lys Glu Lys Tyr Asn Ala Ala
      20      25      30
Val Gln Val Glu Val Val Arg Ala Ser Arg Leu Arg Gln Pro Thr Phe
      35      40      45
Leu Arg Ile Lys Gln Leu Arg Ser Tyr Gln Lys Pro Met Glu Thr Asp
      50      55      60
Leu Val Tyr Ile Glu Lys Ser Pro Asn Tyr Cys Glu Glu Asp Ala Ala
      65      70      75      80
Thr Gly Ser Val Gly Thr Gln Gly Arg Ile Cys Asn Arg Thr Ser Pro
      85      90      95
Gly Ala Asp Gly Cys Asp Thr Met Cys Cys Gly Arg Gly Tyr Asn Thr
      100      105      110
His Gln Tyr Thr Lys Val Trp Gln Cys Asn Cys Lys
      115      120

```

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 365 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met Ala Gly Ser Ala Met Ser Ser Lys Phe Phe Leu Val Ala Leu Ala

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 1 | | | | 5 | | | | 10 | | | | | | 15 | |
| Ile | Phe | Phe | Ser | Phe | Ala | Gln | Val | Val | Ile | Glu | Ala | Asn | Ser | Trp | Trp |
| | | | 20 | | | | | 25 | | | | | 30 | | |
| Ser | Leu | Gly | Met | Asn | Asn | Pro | Val | Gln | Met | Ser | Glu | Val | Tyr | Ile | Ile |
| | | 35 | | | | | 40 | | | | | 45 | | | |
| Gly | Ala | Gln | Pro | Leu | Cys | Ser | Gln | Leu | Ala | Gly | Leu | Ser | Gln | Gly | Gln |
| | 50 | | | | | 55 | | | | | 60 | | | | |
| Lys | Lys | Leu | Cys | His | Leu | Tyr | Gln | Asp | His | Met | Gln | Tyr | Ile | Gly | Glu |
| 65 | | | | 70 | | | | | 75 | | | | | | 80 |
| Gly | Ala | Lys | Thr | Gly | Ile | Lys | Glu | Cys | Gln | Tyr | Gln | Phe | Arg | His | Arg |
| | | | 85 | | | | | 90 | | | | | | 95 | |
| Arg | Trp | Asn | Cys | Ser | Thr | Val | Asp | Asn | Thr | Ser | Val | Phe | Gly | Arg | Val |
| | | | 100 | | | | | 105 | | | | | 110 | | |
| Met | Gln | Ile | Gly | Ser | Arg | Glu | Thr | Ala | Phe | Thr | Tyr | Ala | Val | Ser | Ala |
| | | 115 | | | | | 120 | | | | | 125 | | | |
| Ala | Gly | Val | Val | Asn | Ala | Met | Ser | Arg | Ala | Cys | Arg | Glu | Gly | Glu | Leu |
| | 130 | | | | | 135 | | | | | 140 | | | | |
| Ser | Thr | Cys | Gly | Cys | Ser | Arg | Ala | Ala | Arg | Pro | Lys | Asp | Leu | Pro | Arg |
| 145 | | | | 150 | | | | | 155 | | | | | 160 | |
| Asp | Trp | Leu | Trp | Gly | Gly | Cys | Gly | Asp | Asn | Ile | Asp | Tyr | Gly | Tyr | Arg |
| | | | 165 | | | | | 170 | | | | | | 175 | |
| Phe | Ala | Lys | Glu | Phe | Val | Asp | Ala | Arg | Glu | Arg | Glu | Arg | Ile | His | Ala |
| | | 180 | | | | | 185 | | | | | | 190 | | |
| Lys | Gly | Ser | Tyr | Glu | Ser | Ala | Arg | Ile | Leu | Met | Asn | Leu | His | Asn | Asn |
| | | 195 | | | | | 200 | | | | | 205 | | | |
| Glu | Ala | Gly | Arg | Arg | Thr | Val | Tyr | Asn | Leu | Ala | Asp | Val | Ala | Cys | Lys |
| | 210 | | | | | 215 | | | | | 220 | | | | |
| Cys | His | Gly | Val | Ser | Gly | Ser | Cys | Ser | Leu | Lys | Thr | Cys | Trp | Leu | Gln |
| 225 | | | | 230 | | | | | 235 | | | | | 240 | |
| Leu | Ala | Asp | Phe | Arg | Lys | Val | Gly | Asp | Ala | Leu | Lys | Glu | Lys | Tyr | Asp |
| | | | 245 | | | | | 250 | | | | | | 255 | |
| Ser | Ala | Ala | Ala | Met | Arg | Leu | Asn | Ser | Arg | Gly | Lys | Leu | Val | Gln | Val |
| | | | 260 | | | | 265 | | | | | | 270 | | |
| Asn | Ser | Arg | Phe | Asn | Ser | Pro | Thr | Thr | Gln | Asp | Leu | Val | Tyr | Ile | Asp |
| | | 275 | | | | | 280 | | | | | 285 | | | |
| Pro | Ser | Pro | Asp | Tyr | Cys | Val | Arg | Asn | Glu | Ser | Thr | Gly | Ser | Leu | Gly |
| | 290 | | | | | 295 | | | | | 300 | | | | |
| Thr | Gln | Gly | Arg | Leu | Cys | Asn | Lys | Thr | Ser | Glu | Gly | Met | Asp | Gly | Cys |
| 305 | | | | 310 | | | | | 315 | | | | | 320 | |
| Glu | Leu | Met | Cys | Cys | Gly | Arg | Gly | Tyr | Asp | Gln | Phe | Lys | Thr | Val | Gln |
| | | | 325 | | | | | 330 | | | | | | 335 | |
| Thr | Glu | Arg | Cys | His | Cys | Lys | Phe | His | Trp | Cys | Cys | Tyr | Val | Lys | Cys |
| | | | 340 | | | | 345 | | | | | </ | | | |

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5607 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

ATGTATGTAT GTATGTATGT ATGTATGTAT ACGTGC GTGC ACCTGTGTGT GCTTGGTGTG
60 AGTGGGGCTC AGACATCACC TGATTCCCTG GAACTGGAGT TACAGGTGGC TATAAGCCAC
120

- 42 -

CACTTGGGTG CTGAGAACAG AGTCCGGGCC TCTGGCAGAG CAGTCAGTGC TTTTAGCCAC
180
TGAGCCACTC TCATCCCCCC AATTATGTTC ATCTTGAGTT GGGCAGGTAC GGTGGCGGAA
240
TAGGCCTGTA ATCCAGCAG TCACTGGACC ATCATGGGTT CTACATATTA AACCTTTATG
300
TTAGGTAGGG TCACACAGCA AGATCCGGTC ACAAACCAG CAACAACAAA AACC AAAAGG
360
AGCCAGCTTC TTCCACAAG CATTCTTTCC CTCAGGTCTT CAGCTCCATC TGACAGCTAC
420
TCGGCTGGTG GTCCTATCCT TTCTGAGCCT AGTTGCCAGA GAAACAAGCC CGGTTTCATCT
480
TCATGACTAG CACATCTAAT GATAAGCACA GGTTGACTCA AGGTGCCATA GAGTGACACT
540
AGGTACCCAG AGCGACAGAA TGACACCTAT GAGTGACAGT CGTTAATCAC AAACACACAC
600
ACACACACAC ACACACACAC ACACACACAC TCATGCACCC ACCTGCAAAC ACAATTGCAG
660
CCTTCTGGAC GTCTCCTGTC ACAGCCCCAC CTCCTTCTCTG ATACACTGCG TTAAGTGGTG
720
ACTGTAACAA AATGACTTCA TGCTCTCCCT GTCCTGAGCC AAATTACACA ATTATTTGGA
780
AAGGGCTCAA AATGTTCTTC GTTAGAAGTT TCTGGATACA CCAATACACA GGAGCGTGCA
840
CCCTCAGAAC ACATGTACAC TTTGACTTAA TCTCACGGGT GACACACCGA CGCTTACACT
900
CCCCCTAGCC CACAGAGGCA AACTGCTGGG CGCTTCTGAG TTTCTCACTG CCACCAGCTC
960
GGTTTGCTCA GCCTACCCCC GCACCCCGCG CCCGGGAATC CCTGACCACA GCTCCACCCA
1020
TGCTCTGTCT CCTTCTTTTC CTCTCTGTC CAGCCGTCGG GGTTCCTGGG TGAGGAAGTG
1080
TCTCCACGGA GTCGCTGGCT AGAACCACAA CTTTCATCCT GCCATTGAGA ATAGGGAAGA
1140
GAAGAGACCA CAGCGTAGGG GGGACAGAGG AGACGGACTT CGAGAGGACA GCCCCACCGG
1200
CGCGTGTGGG GGAGGCAATC CAGGCTGCAA ACAGGTTGTC CCCAGCGCAT TGTCCTCGCG
1260
CCCCCTGGCG GATGCTGGTC CCCGACGGGC TCCGGACGCG CAGAAGAGTG AGGCCGGCGC
1320
GCGTGGGAGG CCATCCCAAG GGGAGGGGTC GGCGGCCAGT GCAGACCTGG AGGCCGGGCC
1380
ACCAAGCAGG GGGCGGGGGT GAGCCCCGAC GGTTAGCCTG TCAGCTCTTT GCTCAGACCG
1440
GCAAGAGCCA CAGCTTCGCT CGCCACTCAT TGTCTGTGGC CCTGACCAGT GCGCCCTGGT
1500
GCTTTTAGTG CCGCCCGGGC CCGGAGGGGC AGCCTCTTCT CACTGCAGTC AGCGCCGCAA
1560
CTATAAGAGG CCTATAAGAG GCGGTGCCTC CCGCAGTGGC TGCTTCAGCC CAGCAGCCAG
1620
GACAGCGAAC CATGCTGCCT GCGGCCCCGCC TCCAGACTTA TTAGAGCCAG CCTGGGAACT
1680
CGCATCACTG CCCTCACCGC TGTGTCCAGT CCCACCGTCG CGGACAGCAA CCACAGTCGT
1740
CAGAACCGCA GCACAGAACC AGCAAGGCCA GGCAGGCCAT GGGGCTCTGG GCGCTGCTGC
1800
CCAAGCTGGT TTCTACTACG TTGCTACTGG CACTGACCGC TCTGCCCCGA GCCCTGGCTG
1860
CCAACAGTAG TGGCCGATGG TGGTAAGTGA GCTAGTACGG GGTCCGCCAC TTGTCTGGG
1920
GCAAAGAGCC AGGCACGGGC CTTACCCAGC TCCACGCTG TGGGGATCAC CAACCTACAG
1980

- 43 -

ACCCCCCTCG TGCATTGTGA CTTACATCC AGGGTGCTCA CACCTAGAAC TAGCTCTGCT
2040
GAAGTGGGGC ACATCATTGG CATGCAGAAG CCCAGATACA CCAGGCTCAG AGACCATTCC
2100
CATTTAATAC GACCCCGTTT CTGCTGAGCA ACAGGTCCCA ACCTCGCTGT GGTGGGTGCT
2160
CAGGTGTCCC TTAGGTCTTG AACCAAAAAA AAAAAAAAAA AAAAAAAAAA ACCAGATATT
2220
AGCTTTGAGG TGAGGGAGTG GAATTCCTAA GTTTTTCAAG GTGGGCAAGG CTGCAGGTGG
2280
GGTTTCTCCT CGGGGGCTGA CTTGAAGAAA GGAAGAGCTA AGGTAGCCAT GCCTTTTCTG
2340
TCCACTCACT AGACTCTGGA GCTCAGGGCC AGGCAAGGAT AGGGTGGTAC AGCCTGTATG
2400
GTTAGGATGC AGGTCCCCTC CCCTGGACTG AACCCTTATG CATCCCGCCA GGGGCATCGT
2460
GAACATAGCC TCCTCCACGA ACCTGTTGAC GGATTCCAAG AGTCTGCAGC TGGTGCTCGA
2520
GCCCAGTCTG CAGCTGCTGA GCCGCAAGCA GCGGCGACTG ATCCGACAGA ACCCGGGGAT
2580
CCTGCACAGC GTGAGTGAGG GGCTCCAGAG CGCTGTGCGA GAGTGCAAAT GGCAATTCCG
2640
AAACCGCCGC TGGAAGTGCC CCACTGCTCC GGGGCCCCAC CTCTTCGGCA AGATCGTCAA
2700
CCGAGGTGGG TGCCCAGGAA AGCGACGCTT CCGGGATTAA GGGAAAAGCA GGGTCATCTC
2760
CAGGGCATAG GCGGGCGAAG GCAGGGAAGA CATCCCAGGG TTATATGTGA TCAAAGTGA
2820
AATCGCCTGG TGCCGGCAGT TACCGTAGGT CAGCACCAGA TTCCTTCTAG CCTTGCGTTG
2880
TGAGCATGAT CTTTAACGTT GCTGGCCACT GGCCACAGA AAGGGAATTC CGGATCGTGG
2940
GCGCTGGGCG ACAGCTGTTT TTCCCTAGCC TTCCTCAAAG GTACCTGGGA AGCTGATCTC
3000
TGAGGGCTAG CTAGGGTTGT GCTTCGCACC CAGCAAAGTT TGCCTGCCA ATACTAGTAG
3060
CGATCTTGGC TATGCAGATT TGTCTACTT GGAATCTCC CCTTGGAGCT GCTCTGCTAG
3120
GGCTCTGGAG TCTCAGTAA GCTTAGAGAG GAGGGCATT CATGCTTCGC ACACATGACT
3180
CCAAGGATGT TGGACTGTAG GGTACCAAGT CTTCCAAACA GGGTGCTGAG TTGGCCCCAC
3240
GCCTTCTCTC AACTGATGCG GGGTCGCTTC ACCCACAGGC TGCCGAGAAA CAGCGTTCAT
3300
CTTCGCAATC ACCTCCGCCG GGGTCACACA TTCCGTGGCG CGCTCCTGCT CCGAAGGCTC
3360
CATCGAGTCC TGCACCTGCG ACTACGGCG GCGCGGCCCT GGGGGCCCCG ACTGGCACTG
3420
GGGGGGCTGC AGTGACAACA TCGATTTTGG TCGCTCTTT GGCCGAGAGT TCGTGGAATC
3480
CGGGGAGAAG GGGCGGGACC TACGCTTCCT CATGAACCTT CACAACAACG AGGCAGGGCG
3540
AACGGTACGT CGGTGTGTCC GGAACCAATG GCAGGGGAGA TGTAAGACAG GTGCACGGGG
3600
ACAGAGGCAC AGGGAGGGGC TTCCCGAGAG AGTGGGACTC TAGGAGGGAA GACAGAGAAG
3660
AGGTGGTGGT TGAGGGCAA GAGGTTCCTG AGCTGATGAC AGAACAGAAG AGATTAGCAG
3720
GCTATCAACA CGTGGGATGT ATTGAGATGG CTCCATGGCA CACTTTTGAA AGATAAAAGT
3780
GACTTGCTGG CGTGGAGCAG AGTCTGGCCG AATGTCCCTA TCTCAGCGGG CCATTTTGCA
3840

- 44 -

CTTCCTCTCT CCCGAGCTTA GTCACACCTG GACCTTGGCT GAAGTTTCCA CAGCATCGAC
 3900
 GTGACCCGGG TGGGGTGGGG GTGGGAAGT ATGGGTGGTG GTTCGTGGGA TGTGGCTTT
 3960
 GACCTTTTCT TCCCTCCTCC CCTCGTCCCC TCCTCCCCCA GACCGTGTTT TCTGAGATGC
 4020
 GCCAAGAGTG CAAATGCCAC GGGATGTCCG GCTCCTGCAC GGTGCGCACG TGTGGATGC
 4080
 GGCTGCCCCAC GCTGCGCGCT GTGGGCGACG TGCTGCGCGA CCGCTTCGAC GGCGCCTCCC
 4140
 GCGTCCTTTA CGGCAACCGA GGCAGCAACC GCGCCTCGCG GCGGAGCTG CTGCGCCTGG
 4200
 AGCCCGAAGA CCCC GCGCAC AAGCCTCCCT CCCCTCACGA CCTCGTCTAC TTCGAGAAAT
 4260
 CGCCCAACTT CTGCACGTAC AGTGGCCGCC TGGGCACAGC TGGCACAGCT GGACGAGCTT
 4320
 GCAACAGCTC GTCTCCCGCG CTGGACGGCT GTGAGCTGCT GTGCTGTGGC CGAGGCCACC
 4380
 GCACGCGCAC GCAGCGCGTC ACGGAGCGCT GCAACTGCAC CTTCCACTGG TGCTGCCACG
 4440
 TCAGCTGCCG CAACTGCACG CACACGCGCG TTCTGCACGA GTGTCTATGA GGTGCCGCGC
 4500
 CTCCGGGAAC GGAACGCTC TCTTCCAGTT CTCAGACACA CTCGCTGGTC CTGATGTTTG
 4560
 CCCACCCTAC CGCGTCCAGC CACAGTCCCA GGGTTCATAG CGATCCATCT CTCCCACCTC
 4620
 CTACCTGGGG ACTCCTGAAA CCACTTGCCT GAGTCGGCTC GAACCCTTTT GCCATCCTGA
 4680
 GGGCCCTGAC CCAGCCTACC TCCCTCCCTC TTTGAGGGAG ACTCCTTTTG CACTGCCCCC
 4740
 CAATTTGGCC AGAGGGTGAG AGAAAGATTC TTCTTCTGGG GTGGGGGTGG GGAGGTCAAC
 4800
 TCTTGAAGGT GTTGCGGTTC CTGATGTATT TTGCGCTGTG ACCTCTTTGG GTATTATCAC
 4860
 CTTTCCTTGT CTCTCGGGTC CCTATAGGTC CCTTGAGTTC TCTAACCAGC ACCTCTGGGC
 4920
 TTCAAGGCCT TTCCCTCCC ACCTGTAGCT GAAGAGTTTC CGAGTTGAAA GGGCACGGAA
 4980
 AGCTAAGTGG GAAAGGAGGT TGCTGGACCC AGCAGCAAAA CCCTACATTC TCCTTGCTCTC
 5040
 TGCCTCGGAG CCATTGAACA GCTGTGAACC ATGCCTCCCT CAGCCTCCTC CCACCCCTTC
 5100
 CTGTCTGCC TCCTCATCAC TGTGTAAATA ATTTGCACCG AAATGTGGCC GCAGAGCCAC
 5160
 GCGTTCGGTT ATGTAAATAA AACTATTTAT TGTGCTGGGT TCCAGCCTGG GTTGCAGAGA
 5220
 CCACCTCAC CCCACCTCAC TGCTCCTCTG TTCTGCTCGC CAGTCCTTTT GTTATCCGAC
 5280
 CTTTTTCTC TTTTACCCAG CTCTCATAG GCGCCCTTGC CCACCGGATC AGTATTTCTT
 5340
 TCCACTGTAG CTATTAGTGG CTCCTCGCCC CCACCAATGT AGTATCTTCC TCTGAGGAAT
 5400
 AAAATATCTA TTTTATCAA CGACTCTGGT CCTTGAATCC AGAACACAGC ATGGCTTCCA
 5460
 ACGTCCTCTT CCCTTCCAAT GGAATTGCTT CTCTTCTCAT AGCCAAACAA AAGAGATAGA
 5520
 GTTGTGAAG ATCTCTTTTC CAGGGCCTGA GCAAGGACCC TGAGATCCTG ACCCTTGAT
 5580
 GACCCTAAAT GAGACCAACT AGGGATC
 5607

(2) INFORMATION FOR SEQ ID NO:8:

- 45 -

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2301 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

```
AGCAGAGCGG ACGGGCGCGC GGGAGGCGCG CAGAGCTTTC GGGCTGCAGG CGCTCGCTGC
60
CGCTGGGGAA TTGGGCTGTG GCGGAGGCGG TCCGGGCTGG CCTTTATCGC TCGCTGGGCC
120
CATCGTTTGA AACTTTATCA GCGAGTCGCC ACTCGTCGCA GGACCGAGCG GGGGGCGGGG
180
GCGCGGCGAG GCGGCGGCCG TGACGAGGCG CTCCCGGAGC TGAGCGCTTC TGCTCTGGGC
240
ACGCATGGCG CCCGCACACG GAGTCTGACC TGATGCAGAC GCAAGGGGGT TAATATGAAC
300
GCCCCCTCTG GTGGAATCTG GCTCTGGCTC CCTCTGCTCT TGACCTGGCT CACCCCCGAG
360
GTCAACTCTT CATGGTGGTA CATGAGAGCT ACAGGTGGCT CCTCCAGGGT GATGTGCGAT
420
AATGTGCCAG GCCTGGTGAG CAGCCAGCGG CAGCTGTGTC ACCGACATCC AGATGTGATG
480
CGTGCCATTA GCCAGGGCGT GGCCGAGTGG ACAGCAGAAT GCCAGCACCA GTTCCGCCAG
540
CACCGCTGGA ATTGCAACAC CCTGGACAGG GATCACAGCC TTTTGGCAG GGTCCCTACTC
600
CGAAGTAGTC GGAATCTGC CTTTGTTTAT GCCATCTCCT CAGCTGGAGT TGTATTTGCC
660
ATCACCAGGG CCTGTAGCCA AGGAGAAGTA AAATCCTGTT CCTGTGATCC AAAGAAGATG
720
GGAAGCGCCA AGGACAGCAA AGGCATTTTT GATTGGGGTG GCTGCAGTGA TAACATTGAC
780
TATGGGATCA AATTTGCCCC CGCATTTGTG GATGCAAAGG AAAGGAAAGG AAAGGATGCC
840
AGAGCCCTGA TGAATCTTCA CAACAACAGA GCTGGCAGGA AGGCTGTAAA GCGGTTCTTG
900
AAACAAGAGT GCAAGTGCCA CGGGGTGAGC GGCTCATGTA CTCTCAGGAC ATGCTGGCTG
960
GCCATGGCCG ACTTCAGGAA AACGGGCGAT TATCTCTGGA GGAAGTACAA TGGGGCCATC
1020
CAGGTGGTCA TGAACCAGGA TGGCACAGGT TTCACTGTGG CTAACGAGAG GTTTAAGAAG
1080
CCAACGAAAA ATGACCTCGT GTATTTTGAG AATTCTCCAG ACTACTGTAT CAGGGACCGA
1140
GAGGCAGGCT CCCTGGGTAC AGCAGGCCGT GTGTGCAACC TGACTTCCCG GGCATGGAC
1200
AGCTGTGAAG TCATGTGCTG TGGGAGAGGC TACGACACCT CCCATGTCAC CCGGATGACC
1260
AAGTGTTGGT GTAAGTTCCA CTGGTGCTGC GCCGTGCGCT GTCAGGACTG CCTGGAAGCT
1320
CTGGATGTGC ACACATGCAA GGCCCCAAG AACGCTGACT GGACAACCGC TACATGACCC
1380
CAGCAGGCGT CACCATCCAC CTTCCCTTCT ACAAGGACTC CATTGGATCT GCAAGAACAC
1440
TGGACCTTTG GGTTCCTTCT GGGGGGATAT TTCCTAAGGC ATGTGGCCTT TATCTCAACG
1500
GAAGCCCCCT CTTCCCTCCCT GGGGGCCCCA GGATGGGGGG CCACACGCTG CACCTAAAGC
1560
```

- 46 -

CTACCCTATT CTATCCATCT CCTGGTGTTC TGCAGTCATC TCCCCTCCTG GCGAGTTCTC
 1620
 TTTGGAAATA GCATGACAGG CTGTTTCTAGG GGGAGGGTGG TGGGCCCAGA CCACTGTCTC
 1680
 CACCCACCTT GACGTTTCTT CTTTCTAGAG CAGTTGGCCA AGCAGAAAAA AAAGTGTCTC
 1740
 AAAGGAGCTT TCTCAATGTC TTCCCACAAA TGGTCCCAAT TAAGAAATTC CATACTTCTC
 1800
 TCAGATGGAA CAGTAAAGAA AGCAGAATCA ACTGCCCTG ACTTAACTTT AACTTTTGAA
 1860
 AAGACCAAGA CTTTTGTCTG TACAAGTGGT TTTACAGCTA CCACCCTTAG GGTAATTGGT
 1920
 AATTACCTGG AGAAGAATGG CTTTCAATAC CCTTTTAAGT TTAAAATGTG TATTTTTCAA
 1980
 GGCATTTATT GCCATATTAA AATCTGATGT AACAAGGTGG GGACGTGTGT CCTTTGGTAC
 2040
 TATGGTGTGT TGTATCTTTG TAAGAGCAAA AGCCTCAGAA AGGGATTGCT TTGCATTACT
 2100
 GTCCCCCTGA TATAAAAAAT CTTTAGGGAA TGAGAGTTCC TTCTCACTTA GAATCTGAAG
 2160
 GGAATTAAAA AGAAGATGAA TGGTCTGGCA ATATTCTGTA ACTATTGGGT GAATATGGTG
 2220
 GAAAATAATT TAGTGGATGG AATATCAGAA GTATATCTGT ACAGATCAAG AAAAAAGGA
 2280
 AGAATAAAAT TCCTATATCA T
 2301

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2814 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GAATTCATGT CTTACGGTCA AGGCAGAGGG CCCAGCGCCA CTGCAGCCGC GCCACCTCCC
 60
 AGGGCCGGGC CAGCCCAGGC GTCCGCGCTC TCGGGGTGGA CTCCCCCGGC TGCGCGCTCA
 120
 AGCCGGCGAT GGCTCCTCTC GGATACCTCT TAGTGCTCTG CAGCCTGAAG CAGGCTCTGG
 180
 GCAGCTACCC GATCTGGTGG TCCTTGGCTG TGGGACCCCA GTACTCCTCT CTGAGCACTC
 240
 AGCCCATTCT CTGTGCCAGC ATCCCAGGCC TGGTACCGAA GCAGCTGCGC TTCTGCAGGA
 300
 ACTACGTGGA GATCATGCCC AGCGTGGCTG AGGGTGTCAA AGCGGGCATC CAGGAGTGCC
 360
 AGCACCAATT CCGAGGCCGG CGTTGGAAC TGCACCACCGT CAGCAACAGC CTGGCCATCT
 420
 TTGGCCCTGT TCTGGACAAA GCCACCCGGG AGTCAGCCTT TGTCCATGCC ATCGCCTCCG
 480
 CTGGAGTAGC TTTCGCAGTG ACACGCTCCT GTGCAGAGGG ATCAGCTGCT ATCTGTGGGT
 540
 GCAGCAGCCG CCTCCAGGGC TCCCAGGCG AGGGCTGGAA GTGGGGCGGC TGTAGTGAGG
 600
 ACATTGAATT TGGAGGAATG GTCTCTCGGG AGTTTGCCGA TGCCAGGGAG AACCGGCCGG
 660
 ATGCCCGCTC TGCCATGAAC CGTCACAACA ATGAGGCTGG GCGCCAGGCC ATCGCCAGTC
 720

- 47 -

ACATGCACCT CAAGTGCAAA TGCCACGGGC TATCTGGCAG CTGTGAAGTG AAGACCTGCT
780
GGTGGTCGCA GCCGGACTTC CGCACCATCG GGGATTTCCT CAAGGACAAG TATGACAGTG
840
CCTCGGAGAT GGTGGTAGAG AAACACCGAG AGTCTCGTGG CTGGGTGGAG ACCCTGAGGC
900
CACGTTACAC GTACTTCAAG GTGCCGACAG AACGCGACCT GGTCTACTAC GAGGCCTCAC
960
CCAACTTCTG CGAACCTAAC CCCGAAACCG GCTCCTTCGG GACGCGTGAC CGCACCTGCA
1020
ATGTGAGCTC GCATGGCATA GATGGGTGCG ACCTGTTGTG CTGCGGGCGC GGGCATAACG
1080
CGCGCACTGA GCGACGGAGG GAGAAATGCC ACTGTGTTTT CCATTGGTGC TGCTACGTCA
1140
GCTGCCAGGA GTGCACACGT GTCTATGACG TGCACACCTG CAAGTAGGAG AGCTCCTAAC
1200
ACGGGAGCAG GGTTTCATTCC GAGGGGCAAG GTTCCTACCT GGGGGCGGGG TTCCTACTTG
1260
GAGGGGTCTC TTAATTGGGG ACTCGGTTCT TACTTGAGGG CGGAGATCCT ACCTGTGAGG
1320
GTCTCATACC TAAGGACCCG GTTTCTGCCT TCAGCCTGGG CTCCTATTTG GGATCTGGGT
1380
TCCTTTTTAG GGGAGAAGCT CCTGTCTGGG ATACGGGTTT CTGCCCAGAG GTGGGGCTCC
1440
ACTTGGGGAT GGAATTCCAA TTTGGGCCCG AAGTCCTACC TCAATGGCTT GGAATCCTCT
1500
CTTGACCCGA CAGGGCTCAA ATGGAGACAG GTAAGCTACT CCCTCAACTA GGTGGGGTTC
1560
GTGCGGATGG GTGGGAGGGG AGAGATTAGG GTCCCTCCTC CCAGAGGCAC TGCTCTATCT
1620
AGATACATGA GAGGGTGCTT CAGGGTGGGC CCTATTTGGG CTTGAGGATC CCGTGGGGGC
1680
GGGGCTTCAC CCCGACTGGG TGGAACTTTT GGAGACCCCC TTCCACTGGG GCAAGGCTTC
1740
ACTGAAGACT CATGGGATGG AGCTCCACGG AAGGAGGAGT TCCTGAGCGA GCCTGGGCTC
1800
TGAGCAGGCC ATCCAGCTCC CATCTGGCCC CTTTCCAGTC CTGGTGTAAG GTTCAACCTG
1860
CAAGCCTCAT CTGCGCAGAG CAGGATCTCC TGGCAGAATG AGGCATGGAG AAGAACTCAG
1920
GGGTGATACC AAGACCTAAC AAACCCCGTG CCTGGGTACC TCTTTTAAAG CTCTGCACCC
1980
CTTCTTCAAG GGCTTTCCTA GTCTCCTTGG CAGAGCTTTC CTGAGGAAGA TTTGCAGTCC
2040
CCCAGAGTTC AAGTGAACAC CCATAGAACA GAACAGACTC TATCCTGAGT AGAGAGGGTT
2100
CTCTAGGAAT CTCTATGGGG ACTGCTAGGA AGGATCCTGG GCATGACAGC CTCGTATGAT
2160
AGCCTGCATC CGCTCTGACA CTTAATACTC AGATCTCCCG GGAAACCCAG CTCATCCGGT
2220
CCGTGATGTC CATGCCCCAA ATGCCTCAGA GATGTTGCCT CACTTTGAGT TGTATGAACT
2280
TCGGAGACAT GGGGACACAG TCAAGCCGCA GAGCCAGGGT TGTTTCAGGA CCCATCTGAT
2340
TCCCCAGAGC CTGCTGTTGA GGCAATGGTC ACCAGATCCG TTGGCCACCA CCCTGTCCCG
2400
AGCTTCTCTA GTGTCTGTCT GGCCTGGAAG TGAGGTGCTA CATAAGCCCC ATCTGCCACA
2460
AGAGCTTCCT GATTGGTACC ACTGTGAACC GTCCCTCCCC CTCCAGACAG GGGAGGGGAT
2520
GTGGCCATAC AGGAGTGTGC CCGGAGAGCG CGGAAAGAGG AAGAGAGGCT GCACACGCGT
2580

- 48 -

GGTGACTGAC TGTCTTCTGC CTGGAAC TTT GCGTTCGCGC TTGTAAC TTT ATTTTCAATG
 2640
 CTGCTATATC CACCCACCAC TGGATTTAGA CAAAAGTGAT TTTCTTTTTT TTTTTTCTT
 2700
 TTCTTTCTAT GAAAGAAATT ATTTTAGTTT ATAGTATGTT TGTTTCAAAT AATGGGGAAA
 2760
 GTAAAAAGAG AGAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAA
 2814

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 333 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

| | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| Cys | Lys | Cys | His | Gly | Leu | Ser | Gly | Ser | Cys | Glu | Val | Lys | Thr | Cys | Trp |
| 1 | | | | 5 | | | | | 10 | | | | | 15 | |
| Trp | Ser | Gln | Pro | Asp | Phe | Arg | Ala | Ile | Gly | Asp | Phe | Leu | Lys | Asp | Lys |
| | | | 20 | | | | | 25 | | | | | 30 | | |
| Tyr | Asp | Ser | Ala | Ser | Glu | Met | Val | Val | Glu | Lys | His | Arg | Glu | Ser | Arg |
| | | | 35 | | | | 40 | | | | | 45 | | | |
| Gly | Trp | Val | Glu | Thr | Leu | Arg | Pro | Arg | Tyr | Thr | Tyr | Phe | Lys | Val | Pro |
| | 50 | | | | | 55 | | | | | 60 | | | | |
| Thr | Glu | Arg | Asp | Leu | Val | Tyr | Tyr | Glu | Ala | Ser | Pro | Asn | Phe | Cys | Glu |
| | 65 | | | | 70 | | | | | 75 | | | | 80 | |
| Pro | Asn | Pro | Glu | Thr | Gly | Ser | Phe | Gly | Thr | Arg | Asp | Arg | Thr | Cys | Ans |
| | | | 85 | | | | | 90 | | | | | | 95 | |
| Val | Ser | Ser | His | Gly | Ile | Asp | Gly | Cys | Asp | Leu | Leu | Cys | Cys | Gly | Arg |
| | | | 100 | | | | 105 | | | | | 110 | | | |
| Gly | His | Asn | Ala | Arg | Ala | Glu | Arg | Arg | Glu | Lys | Cys | Arg | Cys | Val | |
| | | 115 | | | | 120 | | | | | 125 | | | | |
| Phe | His | Trp | Cys | Cys | | | | | | | | | | | |
| | | | 130 | | | | | | | | | | | | |

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 399 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

TGTAAGTGCC ACGGGCTGTC GGGCAGCTGC GAGGTGAAGA CATGCTGGTG GTCGCAACCC
 60
 GACTTCCGCG CCATCGGTGA CTTCTCAAG GACAAGTACG ACAGCGCCTC GGAGATGGTG
 120
 GTGGAGAAGC ACCGGGAGTC CCGCGGCTGG GTGGAGACCC TCGCGCCGCG CTACACCTAC
 180
 TTCAAGGTGC CCACGGAGCG CGACCTGGTC TACTACGAGG CCTCGCCCAA CTTCTGCGAG
 240
 CCCAACCTG AGACGGGCTC CTTGGGCACG CGCGACCGCA CCTGCAACGT CAGCTCGCAC
 300

- 49 -

GGCATCGACG GCTGCGACCT GCTGTGCTGC GGCCGCGGCC ACAACGCGCG AGCGGAGCGG
360
CGCCGGGAGA AGTGCCGCTG CGTGTTTCAC TGGTGCTGT
399

- 50 -

What is claimed is:

1. An enriched population of mammalian neural precursor cells committed to a cell fate, said cells being characterized in that they exhibit a stem cell phenotype in the presence of a Wnt polypeptide but not in the absence of said Wnt polypeptide.
2. An enriched population of mammalian dopaminergic neuron precursor cells, said cells being characterized in that they exhibit a stem cell phenotype in the presence of a Wnt polypeptide and differentiate into dopaminergic neurons in the absence of said Wnt polypeptide.
3. The population of claim 2, wherein said Wnt polypeptide is a Wnt-1 class polypeptide.
4. The population of claim 3, wherein said Wnt polypeptide is selected from the group consisting of Wnt-1, Wnt-2, Wnt-3a, Wnt-7a, and Wnt-7b.
5. The population of claim 4, wherein said Wnt polypeptide is Wnt-1.
6. The population of claim 5, wherein said Wnt-1 polypeptide has a sequence that is at least 80% identical to SEQ ID NO: (human Wnt-1).
7. The population of claim 2, wherein said cells are human cells.
8. The population of claim 7, wherein said cells are fetal human cells.
9. The population of claim 2, wherein said cells are porcine cells.

- 51 -

10. An enriched population of mammalian dorsal hindbrain precursor cells, said cells being characterized in that they exhibit a stem cell phenotype in the presence of both a Wnt-1 polypeptide and a Wnt-3a polypeptide but not in
5 the absence of said Wnt-1 polypeptide and said Wnt-3a polypeptide.

11. An enriched population of mammalian hippocampal neuron precursor cells, said cells being characterized in that they exhibit a stem cell phenotype in the presence of a
10 Wnt-3a polypeptide and differentiate into hippocampal neurons in the absence of said Wnt-3a polypeptide..

12. The population of claim 11, wherein said Wnt-3a polypeptide has a sequence that is at least 80% identical to SEQ ID NO: (mouse Wnt-3a).

13. The population of claim 11, wherein said cells
15 are human cells.

14. A method of treating a heterogeneous population of neural cell precursor cells to enrich for dorsal neural precursor cells, comprising culturing said population with
20 Wnt polypeptide, wherein said dorsal neural precursor cells selectively proliferate in the presence of said Wnt polypeptide.

15. A method of stimulating cell proliferation of a dorsal neural precursor cell comprising contacting said cell
25 with a Wnt-1 polypeptide or a Wnt-3a polypeptide.

16. The method of claim 15, wherein said cell is contacted with both a Wnt-1 polypeptide and a Wnt-3a polypeptide.

- 52 -

17. A method of inducing neuronal regeneration in an adult mammal suffering from a neurodegenerative disorder, comprising transplanting into said mammal an enriched population of dorsal neural precursor cells.

5 18. The method of claim 17, wherein said disorder is Parkinson's Disease, Amyotrophic Lateral Sclerosis, Diffuse Lewy Body Disease, Cortical-basal Ganglionic Degeneration, Hallervorden-Spatz Disease, or Myoclonic Epilepsy.

10 19. The method of claim 17, further comprising
/ administering to said mammal a Wnt polypeptide or Wnt agonist.

 20. A method of treating Parkinson's disease, comprising transplanting into the brain of a patient an
15 enriched population of dopaminergic neuron precursor cells.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/08716

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12N 5/06, 5/08

US CL : 435/325, 368, 377

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/325, 368, 377

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, BIOSIS, MEDLINE

search terms: neural, precursor#, progenitor, stem, cell#, human, porcine, wnt#, dorsal, hippocam##, hindbrain, dopamin?

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|---------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|
| X --- Y | US 5,411,883 A (BOSS et al) 02 May 1995, columns 3, 7, 11-13, 17 and 19-20. | 1-9 --- 10-13 |
| X --- A | US 5,589,376 A (ANDERSON et al) 31 December 1996, columns 3-4, 8-9, 11, 13-14 and 16-17. | 1 --- 2-13 |
| X | MOYER et al. Culture, Expansion, and Transplantation of Human Fetal Neural Progenitor Cells. Transplantation Proceedings. June 1997, Vol. 29, No. 4, pages 2040-2041, see entire document. | 1-8, 10-13 |
| X | US 5,656,481 A (BAETGE et al) 12 August 1997, column 30, lines 46-57. | 1, 11-13 |

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

| | |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| * Special categories of cited documents: | *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention |
| *A* document defining the general state of the art which is not considered to be of particular relevance | *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone |
| *E* earlier document published on or after the international filing date | *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art |
| *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) | *G* document member of the same patent family |
| *O* document referring to an oral disclosure, use, exhibition or other means | |
| *P* document published prior to the international filing date but later than the priority date claimed | |

Date of the actual completion of the international search

30 JULY 1998

Date of mailing of the international search report

31 AUG 1998


Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

ROBERT C. HAYES, PH.D.

Telephone No. (703) 308-0196



INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/08716

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-13

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/08716

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s) 1-13, drawn to a population of mammalian neural precursor cells committed to a cell fate.

Group II, claim(s) 14-16, drawn to a method of stimulating proliferation of a heterogeneous population of neural cell precursor cells to enrich for dorsal neural cells.

Group III, claim(s) 17-18 and 20, drawn to a method of inducing neuronal regeneration in an adult mammal comprising transplanting dorsal neural precursor cells.

Group IV, claim(s) 19, drawn to a method of inducing neuronal regeneration in an adult mammal comprising administering a Wnt polypeptide or Wnt agonist.

The inventions listed as Groups I-IV do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

Group I is directed to a population of mammalian neural precursor cells, which is the first product. However, because Boss et al teach an enriched population of porcine or human neuron progenitor cells (i.e., mammalian neural precursor cells), no special technical feature exists for Group I as defined by PCT RULE 13.2, because it does not define a contribution over the prior art. The technical features of Groups II-IV are drawn to methods having different goals, method steps and starting materials, which do not share the same or a corresponding technical feature. Note that PCT Rule 13 does not provide for multiple products or methods within a single application. Because the technical feature of Group I is not a special technical feature, and because the technical features of the Group II-IV inventions are not present in the Group I claims, unity of invention is lacking.